

Combined effects of short-term ocean acidification and heat shock in a benthic copepod *Tigriopus japonicus* Mori

Wei Li^{1,2} · Guodong Han¹ · Yunwei Dong¹ · Atsushi Ishimatsu³ · Bayden D. Russell⁴ · Kunshan Gao¹

Received: 20 March 2015 / Accepted: 6 August 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract Warming of the world's oceans is predicted to have many negative effects on organisms as they have optimal thermal windows. In coastal waters, however, both temperatures and pCO₂ (pH) exhibit diel variations, and biological performances are likely to be modulated by physical and chemical environmental changes. To understand how coastal zooplankton respond to the combined impacts of heat shock and increased pCO₂, the benthic copepod *Tigriopus japonicus* were treated at temperatures of 24, 28, 32 and 36 °C to simulate natural coastal temperatures experienced in warming events, when acclimated in the short term to either ambient (LC, 390 μatm) or future CO₂ (HC, 1000 μatm). HC and heat shock did not induce any mortality of *T. japonicus*, though respiration increased up to 32 °C before being depressed at 36 °C. Feeding rate peaked at 28 °C but did not differ between CO₂ treatments. Expression of heat shock proteins (*hsps* mRNA) was positively related to temperature, with no significant differences between the CO₂ concentrations. Nauplii production was

not affected across all treatments. Our results demonstrate that *T. japonicus* responds more sensitively to heat shocks rather than to seawater acidification; however, ocean acidification may synergistically act with ocean warming to mediate the energy allocation of copepods.

Introduction

Increasing global temperatures are broadly predicted to cause changes to the distributions of species. While virtually all biological processes are affected by temperature, there is recent recognition that the degree to which species are affected will be determined in part by their inherent ability to acclimate and adapt to increased temperatures (Sanford and Kelly 2011). In addition, many habitats demonstrate large variability in environmental conditions, both spatially and temporally, altering predicted biological outcomes (Helmuth et al. 2006; Mislán et al. 2014). Coastal marine environments in particular are usually characterized by large changes in both chemical and physical properties, with diel, seasonal and inter-annual perturbations of temperature, pH and oxygen along with tide cycle, upwelling events and biological activities (Duarte et al. 2013; Marshall et al. 2011; Melzner et al. 2013).

Organisms inhabiting coastal waters experience strong fluctuations of temperature both daily and seasonally (Davison and Pearson 1996). Moreover, sea surface temperature (SST) has been increasing in the last several decades, with shallow coastal waters being strongly impacted (Lima and Wethey 2012), and surface ocean temperature likely to further increase between 0.6 and 2.0 °C by the end of this century (Stocker et al. 2013). Importantly, in addition to this rise in mean temperature, there will be an increase in temperature variation, leading to more extreme

Communicated by H. Pörtner.

Reviewed by undisclosed experts.

✉ Kunshan Gao
ksgao@xmu.edu.cn

- ¹ State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen 361005, China
- ² College of Life and Environmental Sciences, Huangshan University, Huangshan 245041, China
- ³ Institute for East China Sea Research, Nagasaki University, Nagasaki 851-2213, Japan
- ⁴ Swire Institute of Marine Science and School of Biological Sciences, The University of Hong Kong, Hong Kong SAR, China

heating events (Jentsch et al. 2007). For example, Xiamen Bay (Fujian, China) typically has regular, semidiurnal tides (Hu et al. 1988), and water temperatures in the intertidal pools can have a daily temperature range of 25 °C, reaching over 45 °C during low tide in a summer afternoon (Dong et al. personal communication).

Aquatic organisms have different scopes for temperature adaptation and their fitness, and performance can be mediated by the degree to which environmental temperature (and variation) overlaps with their thermal window (Pörtner and Farrell 2008). Though most intertidal animals have developed effective behavioral and physiological adaptations for surviving in this highly variable and harsh environment (Marshall et al. 2011; Somero 2010; Tomanek and Somero 2000), many are still living near their upper thermal limits (Stillman and Somero 2000), meaning that extreme heating events can have deleterious biological effects (Pörtner 2012).

Along with the ocean warming, increasing dissolution of atmospheric CO₂ into seawater is decreasing global average pH in surface oceans, causing ocean acidification (OA), and this acidification is superimposed onto natural fluctuations in pH in coastal waters (Cai et al. 2011; Capone and Hutchins 2013). For example, the surface seawater pH can range from 7.85 to 8.15 in the California upwelling system, and this range will be further lowered to 7.50–7.70 given the continuing and rapid increase in CO₂ emission due to anthropogenic activities (Capone and Hutchins 2013; Sabine et al. 2004). Moreover, eutrophication could contribute additional drop of pH in subsurface of coastal waters, causing more stressful acidification of coastal areas (Cai et al. 2011).

While OA is known to have various effects on processes such as calcification (Comeau et al. 2015; Gao et al. 1993; Ries et al. 2009), other physiological processes in zooplankton such as respiration (Li and Gao 2012; Thomsen and Melzner 2010), pH regulation (Pörtner et al. 2010), feeding (Dupont and Thorndyke 2008; Li and Gao 2012), egg production and nauplius development (Kurihara et al. 2004) were well documented. There were also some studies focus on the combined effects of OA with rising temperature (Munday et al. 2009; Parker et al. 2009). For example, the thermal window of fishes may be narrowed under acidified conditions (Pörtner et al. 2010; Pörtner and Farrell 2008), reducing their thermal scope and fitness with respect to warming. However, what is unknown is how marine organisms, particularly intertidal ectotherms, will respond to rapid and extreme heating events when they are exposed to acidified conditions.

In the scenario of climate change, therefore, it is crucial to investigate the combined effects of reduced pH and heat shock on growth, reproduction and related physiological

performances of intertidal animals. In the present study, we simulated the temperature change observed in natural coastal waters to test the hypothesis that combined effects of OA and extreme heating events increase the costs for somatic maintenance and allocate less energy for growth and reproduction, and show adverse effects on the physiological processes of a benthic copepod *Tigriopus japonicus* Mori. This ecologically important species spreads widely in the coastal waters of western Pacific (China, Japan and Korea) (Jung et al. 2006) and is a common species in the tidal pools on rocky shore where environmental conditions such as temperature and pH demonstrate large daily and seasonal variations (Davenport et al. 1997; McAllen et al. 1999). This species has been extensively studied and is seen as a promising model species for ecotoxicology study given the fully sequenced mitochondrial DNA (Machida et al. 2002; Raisuddin et al. 2007) and can be cultured for several years in indoor conditions (Uye 2005). The species could therefore be taken as a model for studying the relationship between physiological response to temperature fluctuation and pH decline in highly variable coastal areas.

Materials and methods

Experimental setup

This experiment was designed to test the responses of *T. japonicus* to combined heat shock, which they naturally experience in tidal pools during low tide, and the reduced pH under a future OA scenario. It has been shown that pH in intertidal zone could be varied by 0.5 unit during day and night and by more than 1 unit seasonally (Wootton et al. 2008). In the present study, the copepod individuals were acclimated to a short period of different CO₂ treatments (ambient CO₂, LC: 390 µatm, ca. pH 8.15; HC: 1000 µatm, a decrease of ca. 0.4 unit compare with LC) at 20 °C for 24 h, exposed to one of five different temperatures (20, 24, 28, 32 or 36 °C) for 4 h, and returned to 20 °C while being maintained at either LC or HC (see Fig. 1 for experimental design). This heat exposure period was chosen to simulate elevated temperature at low tide in Xiamen Bay. There is a regular semidiurnal tide, which experiences twice of both rising tide and falling tide in 24 h and requires ca. 6 h for a single rising or falling tide, and the surface (1 m layer) seawater temperature of high tide ranges 24.2–27.2 and 12.2–16.4 °C in summer and winter, respectively. Water temperature in intertidal pools during low tides can surpass 45 °C on summer afternoons (Marine Biological Laboratory, Fujian Institute of Oceanology and Division of Marine Biology, Dept. of Biology, Amoy University, 1960). Three different categories of responses to experimental treatments




Fig. 1 Draft of the main design and the timing of each measurement of this study. Expressions below or above the *dark filled rectangle*, *circle* or *up arrows* indicate that specific experiment implement. Copepod treated with LC (390 μatm) and HC (1000 μatm) was given gradually increased, constant and gradually decreased temperature (7 h in total), where the respiration rate was measured after the temperature attain the set value (1.5–5.5 h); *Tigriopus japonicus* acclimated with respective heat shock treatments for 4 h (5.5 h point) were sampled for *hsp 70/hsp 90* mRNA expression test; feeding experiment was carried out for 24 h after 7-h heat shock; female production was measured for 48 h after heat shock treatments; mortality of *T. japonicus* was monitored after 7-h heat shock and the subsequent 72-h culture. Each measurement was an individual experiment, and sample used in one experiment will not be used again

were quantified, physiological (respiration, feeding rate and mortality), molecular (levels of heat shock proteins mRNA) and reproduction (nauplii production).

Adult *T. japonicus* used in this experiment were originally collected from the rocky shore of Xiamen Bay (24°43'N, 118°10'E) and have been cultured in the laboratory in an incubator for 2 years at 20 °C with light intensity of 30 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (12:12 L:D cycle) (a mixed population of male and female with a density of ca. 500 individuals per liter). To maintain optimal conditions, copepods were cultured with offshore seawater (SEATS station, South China Sea, salinity 33 ppt), filtered (0.22 μm), autoclaved at 105 °C and then bubbled with air until pH stabilized at ~8.15. For the entire culture period, the copepods were only fed on the single diatom *Phaeodactylum tricornutum* Bohlin with the initial concentration of $\sim 5 \times 10^4$ cells ml^{-1} .

CO₂ treatments and carbonate chemistry/water conditions

To achieve the different CO₂ treatments, seawater was pre-bubbled with air containing either ambient (390 μatm , LC) or elevated (1000 μatm , HC) CO₂ concentrations using outdoor air and CO₂ chambers (HP1000G-D, Ruihua instrument & equipment Co. Ltd., China), which controls the CO₂ concentration with a variation <3 %. To ensure that treatments were maintained throughout the experiment, pH

of the water was regularly measured with a pH meter (Mettler Toledo DL15 Titrator, Sweden), which was calibrated with NBS buffer solution (Hanna). Parameters of carbonate system (TA, DIC, HCO_3^- , CO_3^{2-} and CO₂) were calculated using the program of CO2SYS for Excel (Lewis and Wallace 1998) with the known temperature, salinity (measured with a handheld refractometer, S/Mill-E, Atago, Japan), nutrients level (seawater forms seats station of Southern China Sea; phosphate is undetectable and can be ignored), pH and $p\text{CO}_2$. The $p\text{CO}_2$ was used for calculation according to the measured $p\text{CO}_2$ in CO₂ chambers with a portable CO₂ detector (Vaisala, CARBOCAP-GM70, Finland). Equilibrium constants for carbonic acid dissociation (K_1 and K_2) were from Roy et al. (1993), and KB for boric acid was after Dickson (1990).

Heat shock treatments

To test the heat shock responses of *T. japonicus*, a rectangular container (length \times width \times height = 75 \times 30 \times 15 cm) made of transparent plastic was divided into five compartments of equal size with plastic dividers. These compartments were used as water baths, with the temperature in each regulated using either hot water or cold water when required to maintain the predetermined temperature. Five target temperatures were set, 20 °C (control), 24, 28, 32 and 36 °C. To simulate natural rates of warming experienced by the copepods on a daily basis, the copepods in bottles with both LC and HC seawater were transferred to 21, 22, 23 and 24 °C in one step, and then, temperature was gradually increased during the next 1.5 h to attain target temperatures. After reaching the target temperature, each water bath was maintained at the designated temperature for 4 h and then gradually decreased back to 20 °C at the same rate as for the heating. The respiration rate was measured after the temperature attains the set value (1.5–5.5 h); *T. japonicus* acclimated with respective heat shock for 4 h (5.5 h point) were sampled for *hsp 70/hsp 90* mRNA expression measurements; feeding experiment was carried out for 24 h after 7-h heat shock; female production was measured for 48 h after heat shock treatments; mortality of *T. japonicus* was monitored after 7-h heat shock and the subsequent 72-h culture (see Fig. 1 for a detailed diagram of the experimental design). Individual copepods were only used for one experiment (i.e., mortality, respiration, feeding, *hsp* mRNA quantification and reproduction measurements, below) so as to not confound other measurements.

Respiration rate and Q_{10}

The respiration rate of adult *T. japonicus* (randomly collected, mixed sex, no gravid female were used) was measured using an oxygen optode (Fibox3, Germany) in sealed

20-ml PerkinElmer glass bottles, during the heat shock treatment. Prior to heating, copepods were placed into bottles containing either LC or HC water ($n = 20$ individuals per bottle; $n = 3$ bottles per treatment) in which an oxygen electrode sensor had been adhered to the inside bottle wall. Oxygen consumption rates were continuously measured over the period that temperature remained stable at the target value (see Fig. 1). No food was added during the respiration trials.

Respiration rates were calculated from the change in oxygen concentration in bottles during the 4-h incubation period:

$$\text{Respiration rate } (\mu\text{g O}_2\text{ind.}^{-1}\text{ h}^{-1}) \\ = (K_{\text{exp.}} - K_{\text{control}}) \times V \times 60/N$$

where “ $K_{\text{exp.}}$ ” and “ K_{control} ” were the oxygen changes per min ($\mu\text{g l}^{-1}\text{ min}^{-1}$) in experimental and control bottles (bottles without copepod), “ V ” the bottle volume (l), “ N ” the copepod number.

The metabolic sensitivity of *T. japonicus* under different pH treatments can be expressed as the Q_{10} , reflecting the change in metabolic rate over a 10 °C temperature gradient. The Q_{10} was calculated according to Atkin and Tjoelker (2003): $Q_{10} = 10^{(\text{slope} \times 10)}$, where slope indicates the regression slope of log10-transformed respiratory rates versus corresponding temperature points (20, 24, 28, 32 and 36 °C).

Feeding rate

The feeding rate was quantified at both LC and HC following the heat shock treatments. As the effects of OA and heat shock may be sustained following removal of the elevated temperature, and because *T. japonicus* primarily feed during the dark (Stearns 1986), the energy consumption (metabolism) and acquisition (feeding) may not temporally coincide (Li and Gao 2012). Therefore, feeding assays were run immediately following return of the water temperature to 20 °C, when copepods were transferred to 110-ml polyethylene bottles containing seawater pre-equilibrated with either LC or HC (as appropriate, $n = 30$ bottles per CO₂ and heat shock combination) at a density of 20 individuals per bottle. *P. tricornutum* were added at a concentration of 5×10^4 cells ml⁻¹. To control for growth of the diatoms over the period of the feeding trials, 20 replicate (each temperature or CO₂ had 2 replicates) bottles containing only diatoms were established. All bottles were then transferred into a dark incubator and held at 20 °C for 24 h, following which feeding and filtering rates were calculated according to Frost (1972) with changes in diatom cell concentrations that measured with a coulter counter (Z2, Beckman Coulter, USA):

$$\text{Feeding rate (cells ind.}^{-1}\text{ h}^{-1}) = V/N \times (\ln C_t - \ln C_{\text{ff}})/t \\ \times (C_{\text{ff}} - C_0)/(\ln C_{\text{ff}} - \ln C_0) \\ \text{Filtering rate (ml ind.}^{-1}\text{ h}^{-1}) = V/N \times (\ln C_t - \ln C_{\text{ff}})/t$$

where “ V ” the volume (ml) of culture, “ N ” the copepod number in each replicate, “ C_0 ” the initial cell concentration, “ C_t ” cell concentration in the control bottle after 24 h (t), “ C_{ff} ” cell concentration in the experimental bottle after 24 h.

Reproductive output

Following the heat shock trial, female *T. japonicus* with egg sacs were randomly selected and transferred into six-well culture plates ($n = 1$ per well, $n = 4$ –6 replicates per treatment) to test the effects of short-term OA and heat shock on nauplii production. Nauplii production requires a period of up to 2 days (Hildebrandt et al. 2014; Kurihara et al. 2004); thus, we checked the nauplii production in the subsequent 2 days after the heat shock to determine the temperature or/and OA effects. During the test, copepod was fed with *P. tricornutum* at a concentration of 5×10^4 cells ml⁻¹ for 2 days under both LC and HC condition at 20 °C. Water in the wells was changed daily and the number of nauplii in the wells enumerated.

Mortality

The mortality of copepods was quantified immediately and for the 3 days following the heat shock trial as described above ($n = 20$ individuals per bottle; $n = 3$ bottles per treatment). Live animals that were not used for other measurements (i.e., reproductive output and heat shock protein production, below) were held at 20 °C in their respective LC or HC conditions and fed with *P. tricornutum* at a concentration of 5×10^4 cells ml⁻¹. The culture water was changed daily with water pre-bubbled to the appropriate CO₂ conditions.

Hsp 70 and hsp 90 measurements

To quantify the production of heat shock proteins under the different treatments, a subsample of *T. japonicus* was taken immediately following the heat shock experiment, washed using phosphate buffer (PBS) for 2–3 times and transferred into 1.5-ml centrifuge tubes containing 0.6 ml PBS (pH 7.80). Specimens were frozen with liquid nitrogen and stored at –80 °C for further processing.

Total RNA was isolated from ca. 50 individuals per treatment ($n = 3$ –7) using Trizol reagent (Invitrogen, Carlsbad, CA, USA). The first strand of cDNA was synthesized using total RNA as a template. Reverse

Table 1 Primer oligonucleotide sequences used for *hsp 70/hsp 90* mRNA expression analysis of *Tigriopus japonicus*

Name	Sequences	References
TJHsp70-RTF	AATTCGCCGACAAGCTCAAG	Rhee et al. (2009)
TJHsp70-RTR	CATACCTCCGGGCATTCCAC	Rhee et al. (2009)
TJHsp90-RTF	CATTCTCAACGCATCCACC	Rhee et al. (2009)
TJHsp90-RTR	AGTTCGGGCATCTCTTCGG	Rhee et al. (2009)
TJ18SF	TCGGGCTGTCTCGTTGGTGATTC	Hwang et al. (2010)
TJ18SR	TGCCACAGTCGACAGTTGATAGG	Hwang et al. (2010)

transcriptase (RT) reactions were performed using Prime-Script™ RT reagent Kit with cDNA Eraser (TAKARA, Shiga, Japan). A partial sequence of the *18S* gene was selected as a reference housekeeping gene to normalize the level of expression. The *18S* gene is a reliable housekeeping gene as previous studies described (Hwang et al. 2010). The levels of *hsp 70* and *hsp 90* expression were quantified using real-time quantitative PCR with primers (Table 1). PCR efficiency of each primer pair was determined by performing standard curves from serial dilutions to ensure that PCR efficiency ranged from 95 to 101 % ($R^2 > 0.99$). PCR was carried out in an ABI 7500 real-time PCR system (Applied Biosystems, MA, USA) in a 20 μ l reaction volume containing 10 μ l of 2 \times Fast-Start DNA Universal SYBR Green Master (Roche, Germany), 0.8 μ l of each primer (10 nmol/ μ l), 1 μ l of cDNA template and 7.4 μ l of RNase-free water. The PCR conditions were as follows: 94 °C/4 min; 35 cycles of 94 °C/30 s, 55 °C/30 s, 72 °C/30 s; and 72 °C/7 min and melt curve confirmation at 95 °C/min, 55 °C/1 min, 80 cycles of 55 °C/10 s with 0.5 °C increase per cycle. All samples were measured in triplicate. Ct (dR) values were analyzed using the ABI 7500 system software (Applied Biosystems, MA, USA). The expression values of *hsp 70* and *hsp 90* mRNA for the various heat treatments were determined as the relative value of *18S* using $2^{-\Delta\Delta CT}$ method for experimental against control treatment (Pfaffl 2001). These experiments were carried out in three technical replicates.

Statistical analysis

Respiration, feeding and filtering rates, reproductive output and HSP production were analyzed using two-way analysis of variance (ANOVA), with both CO₂ concentration (two levels: LC vs. HC) and temperature (five levels: 20, 24, 28, 32 or 36 °C) being fixed and orthogonal ($n = 3$). Where significant effects were detected, pairwise *post hoc* comparison of means was made to determine which factors differed. Q_{10} rate was calculated for the range of temperature treatments and compared between CO₂ treatments using a one-way ANOVA.

Results

Experimental conditions

During the experiment, the carbonate system was significantly different between LC and HC treatments, but stable across all temperature treatments (Table 2). As an example, before and after the respiration experiment the mean pH values in LC and HC were ca. 8.14 and 7.80, respectively (Table 2). HC significantly enhanced the DIC by 9.1–16.2 %, the HCO₃⁻ by 13.6–20.9 % and the CO₂ by 152.3 %, but decreased the CO₃²⁻ by 40.4–48.9 % across all temperature treatments (all $p < 0.05$, Table 2).

Respiration rate, Q_{10} and mortality

No mortality of *T. japonicus* was recorded in either LC or HC during heat shock treatments or in the subsequent 3-day post heat shock culture. For both LC and HC treatments, respiration rates increased with temperature up to 32 °C, but declined under HC and remained steady under LC at 36 °C (Fig. 2a). Respiration rate differed significantly among all temperatures, except between 20 and 24 °C and between 32 and 36 °C ($F_{4,20} = 11.2$, $p < 0.001$, pairwise comparisons). There was also a significant effect of CO₂ concentration on respiration rate (Fig. 2 a, b; $F_{1,20} = 10.7$, $p < 0.005$), with increased respiration of HC animals by 99 % (24 °C) and 60 % (28 °C) in comparison with the LC animals (Fig. 2a). Q_{10} decreased by 30.6 % from LC (2.04 ± 0.29) to HC treatments (1.42 ± 0.22) ($F_{1,4} = 8.8$, $p < 0.05$) (Fig. 2c). There was no significant interactive effect of elevated CO₂ and temperature on respiration ($F_{4,20} = 1.3$, $p > 0.05$) (Table 3).

Feeding and filtering rates

The feeding experiment was carried out for 24 h under constant temperature of 20 °C after the heat shock treatment as described before. Heat shock, but not CO₂, affected both the filtering (temperature, $F_{4,20} = 6.3$, $p < 0.004$; CO₂, $F_{1,20} = 0.004$, $p > 0.9$) and feeding rates (temperature, $F_{4,20} = 6.9$, $p < 0.003$; CO₂, $F_{1,20} = 0.14$, $p > 0.7$) of copepods (Fig. 3 a, b). Both feeding and filtering rates initially

Table 2 Calculated parameters of seawater carbonate system (TA, DIC, HCO_3^- , CO_3^{2-} , CO_2) at the initial and after the respiration experiment

	pH _{NBS}	pCO ₂ (µatm)	TA (µmol Kg ⁻¹)	DIC (µmol Kg ⁻¹)	HCO ₃ ⁻ (µmol Kg ⁻¹)	CO ₃ ²⁻ (µmol Kg ⁻¹)	CO ₂ * (µmol Kg ⁻¹)
<i>Before</i>							
20							
LC	8.14 ± 0.01	399 ± 3.6	2040.6 ± 36.9	1820.5 ± 29.5	1649.0 ± 23.7	158.4 ± 5.9	13.0 ± 0.1
HC	7.83 ± 0.01	1007.0 ± 13.5	2251.3 ± 18.3	2149.9 ± 17.0	2022.6 ± 15.8	94.4 ± 1.6	32.9 ± 0.4
<i>After</i>							
20							
LC	8.12 ± 0.03	399.0 ± 3.6	1984.1 ± 135.4	1765.0 ± 106.4	1598.6 ± 85.9	153.5 ± 20.7	12.9 ± 0.2
HC	7.79 ± 0.01	1007.0 ± 13.5	2126.2 ± 105.4	2036.0 ± 94.1	1918 ± 85.8	85.1 ± 8.8	32.9 ± 0.4
24							
LC	8.14 ± 0.02	399.0 ± 3.6	2041.4 ± 77.5	1821.1 ± 63.4	1649.4 ± 52.0	158.6 ± 11.5	13.0 ± 0.1
HC	7.80 ± 0.03	1007.0 ± 13.5	2072.5 ± 39.6	1987.2 ± 35.2	1873.3 ± 32.1	81.0 ± 3.4	32.9 ± 0.4
28							
LC	8.14 ± 0.01	399.0 ± 3.6	2040.6 ± 36.9	1820.5 ± 29.5	1649 ± 23.7	158.4 ± 5.9	13.0 ± 0.1
HC	7.79 ± 0.01	1007.0 ± 13.5	2125.7 ± 93.1	2035.7 ± 84.0	1917.8 ± 76.9	85.0 ± 7.3	32.9 ± 0.4
32							
LC	8.14 ± 0.01	399.0 ± 3.6	2058.9 ± 51.2	1835.8 ± 42.2	1661.8 ± 34.8	160.9 ± 7.5	13.0 ± 0.1
HC	7.81 ± 0.01	1007.0 ± 13.5	2234.6 ± 111.2	2134.6 ± 99.0	2008.4 ± 89.8	93.3 ± 9.6	32.9 ± 0.4
36							
LC	8.15 ± 0.02	399.0 ± 3.6	2077.5 ± 67.5	1851.3 ± 55.2	1674.8 ± 45.3	163.5 ± 10.0	13.0 ± 0.1
HC	7.83 ± 0.02	1007.0 ± 13.5	2229.4 ± 18.3	2131.0 ± 17.8	2005.7 ± 16.8	92.2 ± 0.8	33.1 ± 0.3

Parameters of total alkalinity (TA), dissolved inorganic carbon (DIC), HCO_3^- , CO_3^{2-} and dissolved aqueous carbon dioxide (CO_2^*) were calculated based on the known value of pH_{NBS}, pCO₂, salinity, nutrient concentration and temperature using the CO2Sys.exe program. Data are the mean ± SD of 3 measurements

showed a positive relationship with temperature (this trend was present from 24 °C for HC group) but reached maximum values at 28 °C, from which they declined to the minimum values at 36 °C (Fig. 3a, b). There was no significant interactive effect of elevated CO₂ and temperature on feeding and filtering rates ($F_{4,20} = 1.3$, $p > 0.05$) (Table 3).

Hsp 70 and hsp 90 mRNA

Hsp 70 and hsp 90 mRNA production showed a positive relationship with temperature rise, but were not affected by elevated CO₂ ($F_{1,32} = 0.45$, $p > 0.5$). Hsp 70 mRNA levels increased from 20 to 24 °C and did not change from 28 to 32 °C, before rapidly increasing to 36 °C (Fig. 4a, $F_{4,32} = 7.7$, $p < 0.001$, pairwise comparisons). Hsp 90 showed the same pattern as hsp 70, except that the increase with temperature was more consistent in the LC treatment than under HC conditions (Fig. 4b, Temp × CO₂ interaction: $F_{4,32} = 2.9$, $p < 0.05$) (Table 3).

Reproductive output

Neither the heat shock nor CO₂ treatments caused any changes in the production rate of nauplii in the 2 days following the heat shock (Table 4; all $p > 0.3$).

Discussion

Warming of the world's oceans is already driving species range shifts and changes to community assemblages (Wernberg et al. 2012). Yet, the underlying mechanisms for these changes are often not well understood. In coastal waters in particular, where daily variation can be greater than seasonal change (Helmuth et al. 2006), biological responses are more likely to be a result of short-term extreme events than changes to average conditions (Helmuth et al. 2014). Here, we show that combined heat shock and OA did not affect the mortality of *T. japonicus*, a copepod which inhabits a highly variable intertidal environment, even though they were cultured under indoor condition for 2 years in this study. The mechanism underlying this resistance is likely to be twofold, according to the present results, the regulation of metabolic rates and the production of protective heat shock proteins.

Below the upper thermal limit, metabolic rates in aquatic invertebrates are predominantly driven by temperature (Perry et al. 2005; Rosenzweig et al. 2008). In addition, temperature variation is known to dramatically mediate physiological processes such as the respiration, enzyme activity, production and development of organisms (Angilletta 2009; Pörtner and Farrell 2008). In face of the extreme temperature, organisms can passively mediate




Fig. 2 Respiration rates (a), the \log_{10} (respiration) (b) and the calculated Q_{10} (c) of *Tigriopus japonicus* treated with different temperatures and the respiration represent the calculated oxygen consumption rate during the 4 h under respective stable temperature conditions. Data are the mean \pm SD of 3 measurements. The “*” above the symbol indicates significant differences between treatments at $p < 0.05$ level

their metabolism, for example, through a reduction in their metabolic rate below the basal level (called dormancy), which is seen as an energy conservation process (Marshall et al. 2011; McAllen et al. 1999). Further, upper thermal tolerance limits differ among species and different populations of the same species from different environments (Stillman and Somero 2000). In particular, populations that inhabit variable environments and are exposed to extremes tend to demonstrate physiological coping mechanisms (Willmer et al. 2005). In the present study, there was no mortality of *T. japonicus* that long term cultured in indoor condition, even in the most extreme temperature and CO_2 treatments, which could be expected as *T. japonicus* inhabit

Table 3 Two-way ANOVA results for female production, respiration, *hsp 70*, *hsp 90*, grazing and filtering rate of CO_2 and temperature on *T. japonicus*

	SS	DF	MS	F	P
<i>Female production</i>					
CO_2	2.842	1	2.842	0.167	0.685
Temperature	92.728	4	23.182	1.362	0.265
CO_2 * temperature	34.018	4	8.504	0.500	0.736
<i>Respiration</i>					
CO_2	0.003	1	0.003	10.720	0.004
Temperature	0.011	4	0.003	11.200	0.000
CO_2 * temperature	0.001	4	0.000	1.312	0.300
<i>Hsp70</i>					
CO_2	35.138	4	8.785	7.744	0.000
Temperature	0.431	1	0.431	0.380	0.542
CO_2 * temperature	0.702	4	0.175	0.155	0.960
<i>Hsp90</i>					
CO_2	73.870	4	18.468	30.765	0.000
Temperature	0.243	1	0.243	0.404	0.529
CO_2 * temperature	6.986	4	1.747	2.910	0.037
<i>Grazing</i>					
CO_2	120,45.453	1	120,45.453	0.148	0.704
Temperature	2,270,912.733	4	567,728.183	6.982	0.001
CO_2 * temperature	320,373.711	4	80,093.428	0.985	0.438
<i>Filtering</i>					
CO_2	0.000	1	0.000	0.004	0.949
Temperature	0.001	4	0.000	6.323	0.002
CO_2 * temperature	0.000	4	0.000	0.851	0.510

the nearshore zone and are known to be a tolerant species with strong adaptive ability (Davenport et al. 1997; Kwok and Leung 2005). The used indoor cultured species which can be considered as representative for copepod long term acclimated under constant condition (type of stable condition) and may helpful to clarify the responses to the sudden heat shocks under OA condition.

Both elevated temperature and $p\text{CO}_2$ affect the metabolic rate of the copepods. The respiration rate of *T. brevicornis* is known to have a linear relationship with temperature increases from 5 to 30 $^{\circ}\text{C}$; however, at temperatures above 35 $^{\circ}\text{C}$ the respiration rate is not stimulated further (McAllen et al. 1999), which is in accordance with our present result. Yet, the effects of CO_2 on respiration of copepods are not consistent among taxa; high CO_2 elevates respiration in *Centropages tenuiremis* (Li and Gao 2012), but it only had a significant effect on respiration of *Acartia clausi* in combination with high temperature (Zervoudaki et al. 2013). In another recent study, elevated temperature but not CO_2 significantly stimulated the respiration of female *Calanus hyperboreus* (Hildebrandt et al. 2014). The different responses of the aerobic performance to OA may




Fig. 3 Feeding (a) and filtering (b) rates of *Tigriopus japonicus* treated with different temperatures for 7 h (1.5-h increase, 4-h constant and 1.5-h decrease) and then transferred back to 20 °C under dark condition for subsequent 24 h. Data are the mean \pm SD of 3 measurements

be related to differential resistance or energy requirements among species, especially under extreme thermal conditions, at which thermal windows will potentially be narrowed (Pörtner and Farrell 2008).

Increasing metabolic rate under future OA conditions may relate to acid–base regulation in many marine animals, which is an energy-demanding process that can be affected by seawater acidification due to the changed pH gradient between intracellular and extracellular compartments (Pörtner et al. 2000). Therefore, it is not surprising that enhanced metabolic rate or Na/K⁺ ATPase activities (an indicator of ion regulatory ability) can be evident in moderate hypercapnic conditions (Melzner et al. 2009; Wood et al. 2008). In the present study, short-term exposure to acidification caused elevated respiration in *T. japonicus*. Yet, *Tigriopus* spp. are known to have excellent osmoregulation and/or hemolymph regulation ability (Davenport et al. 1997), likely because of the coastal areas they inhabit in always with a large natural range of environmental variations (salinity, pCO₂, etc.). It is likely that respiration was stimulated at 24 and 28 °C under OA but not in higher temperatures (32 and 36 °C) because the combined effects caused metabolic down-regulation at a lower temperature (e.g., also demonstrated in *Calanus glacialis*; Hildebrandt et al.




Fig. 4 Relative *hsp 70* (a) and *hsp 90* (b) gene expression of *Tigriopus japonicus* after treated with different temperatures and kept in target temperature for 4 h; data are the mean \pm SD of 3–7 measurements

2014). It is possible, therefore, that the elevated respiration rates seen here are a shock response to the shorter-term exposure to acidified conditions. Indeed, changes to metabolic processes in response to acidification are generally followed by an increase in energy acquisition or metabolic cost (Thor and Oliva 2015; Thomsen and Melzner 2010), a mechanism that has been shown to increase survival in several taxa (Burnell et al. 2013; Tunnicliffe et al. 2009), and the response that is tested could be mediated by food concentration in the same species within different populations (Thor and Oliva 2015).

To avoid mortality as temperature increases, the elevated metabolic demands must be met by increased consumption of food (Burnell et al. 2013). Feeding and filtering rates in *T. japonicus* increased from 20 to 28 °C, but decreased above 28 °C, with the lowest feeding and filtering rates at 36 °C. In contrast, metabolic rate continued to increase to 32 °C, before being suppressed at 36 °C. Therefore, at the highest temperature the copepods had higher metabolic demands than they were able to service with feeding, meaning that they were drawing on energetic reserves. While there was no mortality from this single extreme heating event, subsequent events or sustained extreme warming (e.g., the 5 weeks of warming observed

Table 4 Nauplii production of *Tigriopus japonicus* (ind. $^{-1}$ 2 days $^{-1}$) in the 2 days following heat shock at temperatures of 20–36 °C for 7 h

	Temperature (°C)				
	20	24	28	32	36
LC	15.75 ± 7.37	11.33 ± 3.27	12.60 ± 1.95	13.00 ± 4.69	14.60 ± 3.71
HC	14.20 ± 4.32	14.67 ± 4.27	12.75 ± 5.62	13.50 ± 5.07	10.60 ± 0.89

Following the heat shock, *T. japonicus* were maintained in either their LC or HC treatment, but the temperature was reduced back to 20 °C to simulate being re-immersed in seawater. ($n = 3-6$ individuals per mean)

in Western Australia) (Wernberg et al. 2012) may cause mortality because of cumulative effects and metabolic debt. Indeed, such mismatch between metabolic rates and feeding are becoming increasingly recognized in invertebrates (Lemoine and Burkepille 2012) and may lead to mortality on longer time scales (Mertens et al. in press). Alternatively, the mismatch between energy acquisition (reduced feeding) and consumption (higher respiration) may also be due to the exposure duration to such extreme fluctuations not being sufficient for the copepods to adapt (Li and Gao 2012), but subsequent events combined with the short generation time of these animals may enable them to adapt to these new conditions. For example, acclimated to higher temperatures and OA for longer periods (5 weeks), the intertidal snail *Littorina littorea* show enhanced feeding compared with individuals acclimated for shorter periods (2 weeks) (Russell et al. 2013). Further, enhanced grazing rate of micro-zooplankton as a consequence of OA and enhanced temperature was also observed based on a ca. 20-day mesocosm study (Kim et al. 2010). In these cases, long-term survival under altered conditions seems to be related to increasing consumption of food to compensate for increased metabolic costs.

Heat shock proteins, a defensive strategy that can play a critical protective role when exposed to extreme temperature stress, were strongly up-regulated in response to warming in *T. japonicus* (Raisuddin et al. 2007; Rhee et al. 2009). HSPs are important molecular chaperones which help avoid damage of cells by aiding in the repair of nucleoprotein and in preventing coagulation (Bierkens 2000), potentially improving the adaptive ability of organisms against high temperature and other various environmental stresses (Ohtsuka and Hata 2000). Thus, enhanced expression of HSPs under thermal stress is a common defense strategy (Feder and Hofmann 1999; Lee et al. 2008). Again, the 100 % survival of *T. japonicus* indicates that these copepods are resistant to thermal stress, and it is likely that this is at least partially a result of the protective effects of heat shock protein that evidenced in up-regulated *hsp* 70 and *hsp* 90 mRNA. However, the synthesis and function of heat shock proteins are energy-intensive, meaning that more energy must be allocated into somatic maintenance in high temperature (Sokolova et al. 2012). Whether this extra cost contributes to the increased metabolic rate in elevated temperature is unclear, especially as the highest *hsp* mRNA expression was at 36 °C when metabolic rate was depressed. Regardless, the extra energy that needs to be allocated to metabolic processes and HSP production results in less energy being allocated to growth and reproduction (Sokolova et al. 2012). Thus, under extended or repeated heating events, increased individual survival would come at the expense of population growth through reduced reproductive output (Hofmann and Todgham 2010).

While OA is ongoing with ocean warming, heat shock in intertidal areas or coastal waters could reduce the overall physiological performances (Marshall et al. 2011), thereby strongly influencing population dynamics of rocky intertidal species (Garrabou et al. 2009). Therefore, understanding the combined effects of acidification and warming in the context of natural fluctuation in abiotic conditions will provide insights into the potential influences of global change on species inhabiting coastal environments. For most experimental studies on marine organisms, their growth conditions are usually quite different from that in the sea, since they are often grown in vessels under constant conditions during experimental periods. Therefore, it has to be pointed out that cautions in interpreting the data obtained in enclosures should be taken, especially to reflect ecological or global change impacts. The copepod used in this study was maintained under indoor constant temperature and light for almost 2 years, which may lead to evolutionary changes in its physiological characteristics. Consequently, the data obtained here can only reflect the copepod's conditional responses to future climate changes. To achieve the goal we aimed, further tests in the laboratory and/or in field are needed to address the copepod's responses to multiple stressors. Obviously, multifaceted, but long-term, in situ experiments are needed to shed light on whether species inhabiting marine harsh environments can adapt to, and survive, global changes superimposed on the natural variation in abiotic conditions (Russell et al. 2012).

Acknowledgments This study was supported by National Natural Science Foundation (Nos. 41430967; 41120164007), Joint project of NSFC and Shandong province (Grant No. U1406403), Strategic Priority Research Program of CAS (Grant No. XDA11020302), SOA (GASI-03-01-02-04) and China–Japan collaboration project from MOST (S2012GR0290). YWD is funded by Program for New Century Excellent Talents of Ministry of Education, China. BDR is funded by an Australian Research Council Discovery Grant. WL and BDR are funded by MEL Visiting Fellowship of State Key Laboratory of Marine Environmental Science, Xiamen University. We would like to thank Dr. Guizhong Wang (Xiamen University, China) for providing *Tigriopus japonicus*; we also thank Yahe Li and Nana Liu (Xiamen University, China) for their kind assistance during the experiments.

Compliance with ethical standards

Ethical standard This study used copepod as material. All procedures performed in studies involving copepod were in accordance with the ethical standards of the institution or practice at which the studies were conducted.

Conflict of interest The authors declare that they have no conflict of interest.

References

- Angilletta MJ (2009) Thermal adaptation: a theoretical and empirical synthesis. Oxford University Press, Oxford
- Atkin OK, Tjoelker MG (2003) Thermal acclimation and the dynamic response of plant respiration to temperature. *Trends Plant Sci* 8:343–351
- Bierkens JGEA (2000) Applications and pitfalls of stress-proteins in biomonitoring. *Toxicology* 153:61–72
- Burnell O, Russell B, Irving A, Connell S (2013) Eutrophication offsets increased sea urchin grazing on seagrass caused by ocean warming and acidification. *Mar Ecol Prog Ser* 485:37–46
- Cai W-J et al (2011) Acidification of subsurface coastal waters enhanced by eutrophication. *Nat Geosci* 4:766–770
- Capone DG, Hutchins DA (2013) Microbial biogeochemistry of coastal upwelling regimes in a changing ocean. *Nat Geosci* 6:711–717
- Comeau S, Carpenter R, Lantz C, Edmunds P (2015) Ocean acidification accelerates dissolution of experimental coral reef communities. *Biogeosciences* 12:365–372
- Davenport J, Barnett PRO, McAllen RJ (1997) Environmental tolerances of three species of the harpacticoid copepod genus *Tigriopus*. *J Mar Biol Assoc UK* 77:3–16
- Davison IR, Pearson GA (1996) Stress tolerance in intertidal seaweeds. *J Phycol* 32:197–211
- Dickson AG (1990) Standard potential of the reaction: $\text{AgCl(s)} + \frac{1}{2} \text{H}_2(\text{g}) = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic seawater from 273.15 to 318.15 K. *J Chem Thermodyn* 22:113–127
- Duarte CM et al (2013) Is ocean acidification an open-ocean syndrome? Understanding anthropogenic impacts on seawater pH. *Estuar Coast* 36:221–236
- Dupont S, Thorndyke MC (2008) Ocean acidification and its impact on the early life-history stages of marine animals. In: Briand F (ed) Impacts of acidification on biological, chemical and physical systems in the Mediterranean and Black Seas. CIESM Monographs, Monaco, pp 89–97
- Feder ME, Hofmann GE (1999) Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annu Rev Physiol* 61:243–282
- Frost BW (1972) Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod *Calanus pacificus*. *Limnol Oceanogr* 17:805–815
- Gao K, Aruga Y, Asada K, Ishihara T, Akano T, Kiyohara M (1993) Calcification in the articulated coralline alga *Corallina pilulifera*, with special reference to the effect of elevated CO_2 concentration. *Mar Biol* 117:129–132
- Garrabou J et al (2009) Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. *Glob Change Biol* 15:1090–1103
- Helmuth B et al (2006) Mosaic patterns of thermal stress in the rocky intertidal zone: implications for climate change. *Ecol Monogr* 76:461–479
- Helmuth B et al (2014) Beyond long-term averages: making biological sense of a rapidly changing world. *Clim Change Responses* 1:6
- Hildebrandt N, Niehoff B, Sartoris FJ (2014) Long-term effects of elevated CO_2 and temperature on the Arctic calanoid copepods *Calanus glacialis* and *C. hyperboreus*. *Mar Pollut Bull* 80:59–70
- Hofmann GE, Todgham AE (2010) Living in the now: physiological mechanisms to tolerate a rapidly changing environment. *Annu Rev Physiol* 72:127–145
- Hu J, Fu Z, Wu L (1988) Tidal residual circulation in the regular semi-diurnal tide. *J Xiamen Univ (Nat Sci)* 27:323–327

- Hwang D-S et al (2010) Modulation of *p53* gene expression in the intertidal copepod *Tigriopus japonicus* exposed to alkylphenols. *Mar Environ Res* 69:S77–S80
- Jentsch A, Kreyling J, Beierkuhnlein C (2007) A new generation of climate-change experiments: events, not trends. *Front Ecol Environ* 5:365–374
- Jung S-O et al (2006) The complete mitochondrial genome of the intertidal copepod *Tigriopus* sp. (Copepoda, Harpacticidae) from Korea and phylogenetic considerations. *J Exp Mar Biol Ecol* 333:251–262
- Kim J-M, Lee K, Yang EJ, Shin K, Noh JH, K-t Park, Hyun B, Jeong H-J, Kim J-H, Kim KY (2010) Enhanced production of oceanic dimethylsulfide resulting from CO₂-induced grazing activity in a high CO₂ world. *Environ Sci Technol* 44:8140–8143
- Kurihara H, Shimode S, Shirayama Y (2004) Effects of raised CO₂ concentration on the egg production rate and early development of two marine copepods (*Acartia steueri* and *Acartia erythraea*). *Mar Pollut Bull* 49:721–727
- Kwok KWH, Leung KMY (2005) Toxicity of antifouling biocides to the intertidal harpacticoid copepod *Tigriopus japonicus* (Crustacea, Copepoda): effects of temperature and salinity. *Mar Pollut Bull* 51:830–837
- Lee K-W, Raisuddin S, Hwang D-S, Park HG, Dahms H-U, Ahn I-Y, Lee JS (2008) Two-generation toxicity study on the copepod model species *Tigriopus japonicus*. *Chemosphere* 72:1359–1365
- Lemoine NP, Burkepille DE (2012) Temperature-induced mismatches between consumption and metabolism reduce consumer fitness. *Ecology* 93:2483–2489
- Lewis E, Wallace DWR (1998) Program developed for CO₂ system calculations. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee
- Li W, Gao K (2012) A marine secondary producer respire and feeds more in a high CO₂ ocean. *Mar Pollut Bull* 64:699–703
- Lima FP, Wethey DS (2012) Three decades of high-resolution coastal sea surface temperatures reveal more than warming. *Nat Commun* 3:704
- Machida RJ, Miya MU, Nishida M, Nishida S (2002) Complete mitochondrial DNA sequence of *Tigriopus japonicus* (Crustacea: Copepoda). *Mar Biotechnol* 4:406–417
- Marine Biological Laboratory, Fujian Institute of Oceanology, Division of Marine Biology, Dept. of Biology, Amoy University (1960) Studies on the littoral ecology of Amoy and its vicinity. *J Xiamen Univ (Nat Sci)* 3:74–95
- Marshall DJ, Dong YW, McQuaid CD, Williams GA (2011) Thermal adaptation in the intertidal snail *Echinolittorina malaccana* contradicts current theory by revealing the crucial roles of resting metabolism. *J Exp Biol* 214:3649–3657
- McAllen R, Taylor AC, Davenport J (1999) The effects of temperature and oxygen partial pressure on the rate of oxygen consumption of the high-shore rock pool copepod *Tigriopus brevicornis*. *Comp Biochem Physiol A Mol Integr Physiol* 123:195–202
- Melzner F, Göbel S, Langenbuch M, Gutowska MA, Pörtner H-O, Lucassen M (2009) Swimming performance in Atlantic Cod (*Gadus morhua*) following long-term (4–12 months) acclimation to elevated seawater PCO₂. *Aquat Toxicol* 92:30–37
- Melzner F et al (2013) Future ocean acidification will be amplified by hypoxia in coastal habitats. *Mar Biol* 160:1875–1888
- Mislan K, Helmuth B, Wethey DS (2014) Geographical variation in climatic sensitivity of intertidal mussel zonation. *Global Ecol Biogeogr* 23:744–756
- Munday PL, Dixson DL, Donelson JM, Jones GP, Pratchett MS, Devitsina GV, Døving KB (2009) Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proc Natl Acad Sci USA* 106:1848–1852
- Ohtsuka K, Hata M (2000) Molecular chaperone function of mammalian *Hsp 70* and *Hsp 40*-a review. *Int J Hyperth* 16:231–245
- Parker LM, Ross PM, O'Connor WA (2009) The effect of ocean acidification and temperature on the fertilization and embryonic development of the Sydney rock oyster *Saccostrea glomerata* (Gould 1850). *Glob Chang Biol* 15:2123–2136
- Perry AL, Low PJ, Ellis JR, Reynolds JD (2005) Climate change and distribution shifts in marine fishes. *Science* 308:1912–1915
- Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29:e45
- Pörtner HO (2012) Integrating climate-related stressor effects on marine organisms: unifying principles linking molecule to ecosystem-level changes. *Mar Ecol Prog Ser* 470:273–290
- Pörtner HO, Farrell AP (2008) Physiology and climate change. *Science* 322:690–692
- Pörtner HO, Bock C, Reipschläger A (2000) Modulation of the cost of p_{H_i} regulation during metabolic depression: a ³¹P-NMR study in invertebrate (*Sipunculus nudus*) isolated muscle. *J Exp Biol* 203:2417–2428
- Pörtner HO, Dupont S, Melzner F, Storch D, Thorndyke M (2010) Studies of metabolic rate and other characters across life stages. In: Riebesell U, Fabry VJ, Hansson L, Gattuso J-P (eds) Guide to best practices ocean acidification and data reporting. Publications Office of the European Union, Luxembourg, pp 137–165
- Raisuddin S, Kwok KW, Leung KM, Schlenk D, Lee JS (2007) The copepod *Tigriopus*: a promising marine model organism for ecotoxicology and environmental genomics. *Aquat Toxicol* 83:161–173
- Rhee J-S et al (2009) Heat shock protein (*Hsp*) gene responses of the intertidal copepod *Tigriopus japonicus* to environmental toxicants. *Comp Biochem Physiol C Toxicol Pharmacol* 149:104–112
- Ries JB, Cohen AL, McCorkle DC (2009) Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification. *Geology* 37:1131–1134
- Rosenzweig C et al (2008) Attributing physical and biological impacts to anthropogenic climate change. *Nature* 453:353–357
- Roy RN et al (1993) The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45 C. *Mar Chem* 44:249–267
- Russell BD, Harley CD, Wernberg T, Mieszkowska N, Widdicombe S, Hall-Spencer JM, Connell SD (2012) Predicting ecosystem shifts requires new approaches that integrate the effects of climate change across entire systems. *Biol Lett* 8:164–166
- Russell BD, Connell SD, Findlay HS, Tait K, Widdicombe S, Mieszkowska N (2013) Ocean acidification and rising temperatures may increase biofilm primary productivity but decrease grazer consumption. *Philos Trans R Soc Lond B Biol Sci* 368:20120438
- Sabine CL et al (2004) The oceanic sink for anthropogenic CO₂. *Science* 305:367–371
- Sanford E, Kelly MW (2011) Local adaptation in marine invertebrates. *Annu Rev Mar Sci* 3:509–535
- Sokolova IM, Frederich M, Bagwe R, Lannig G, Sukhotin AA (2012) Energy homeostasis as an integrative tool for assessing limits of environmental stress tolerance in aquatic invertebrates. *Mar Environ Res* 79:1–15
- Somero GN (2010) The physiology of climate change: how potentials for acclimatization and genetic adaptation will determine 'winners' and 'losers'. *J Exp Biol* 213:912–920
- Stearns DE (1986) Copepod grazing behavior in simulated natural light and its relation to nocturnal feeding. *Mar Ecol Prog Ser* 30:65–76
- Stillman JH, Somero GN (2000) A comparative analysis of the upper thermal tolerance limits of eastern Pacific porcelain crabs, genus *Petrolisthes*: influences of latitude, vertical zonation, acclimation, and phylogeny. *Physiol Biochem Zool* 73:200–208
- Stocker TF et al (2013) Climate change 2013: the physical science basis. In: Intergovernmental panel on climate change, working

- group I contribution to the IPCC Fifth Assessment Report (AR5). Cambridge University Press, New York
- Thomsen J, Melzner F (2010) Moderate seawater acidification does not elicit long-term metabolic depression in the blue mussel *Mytilus edulis*. *Mar Biol* 157:2667–2676
- Thor P, Oliva EO (2015) Ocean acidification elicits different energetic responses in an Arctic and a boreal population of the copepod *Pseudocalanus acuspes*. *Mar Biol* 162:799–807
- Tomanek L, Somero GN (2000) Time course and magnitude of synthesis of heat-shock proteins in congeneric marine snails (Genus *Tegula*) from different tidal heights. *Physiol Biochem Zool* 73:249–256
- Tunncliffe V, Davies KT, Butterfield DA, Embley RW, Rose JM, Chadwick WW Jr (2009) Survival of mussels in extremely acidic waters on a submarine volcano. *Nat Geosci* 2:344–348
- Uye S (2005) A brief review of mass culture of copepods used for fish food in Japanese mariculture and a proposed plan to use high biomass natural populations of brackish-water copepods. In: Lee C, O'Bryen PJ, Marcus NH (eds) *Copepods in aquaculture*. Blackwell Publishing, Oxford, pp 75–90
- Wernberg T, Smale DA, Thomsen MS (2012) A decade of climate change experiments on marine organisms: procedures, patterns and problems. *Glob Change Biol* 18:1491–1498
- Willmer P, Stone G, Johnston I (2005) *Environmental physiology of animals*. Blackwell, Malden
- Wood HL, Spicer JI, Widdicombe S (2008) Ocean acidification may increase calcification rates, but at a cost. *Proc R Soc Lond B Biol Sci* 275:1767–1773
- Wootton JT, Pfister CA, Forester JD (2008) Dynamic patterns and ecological impacts of declining ocean pH in a high-resolution multi-year dataset. *Proc Natl Acad Sci USA* 105:18848–18853
- Zervoudaki S, Frangoulis C, Giannoudi L, Krasakopoulou E (2013) Effects of low pH and raised temperature on egg production, hatching and metabolic rates of a Mediterranean copepod species (*Acartia clausi*) under oligotrophic conditions. *Mediterr Mar Sci* 15:74–83