



Short communication

Proteomic analysis by iTRAQ in red claw crayfish, *Cherax quadricarinatus*, hematopoietic tissue cells post white spot syndrome virus infection



Joseph Jeswin^{a,1}, Xiao-lu Xie^{a,1}, Qiao-lin Ji^a, Ke-jian Wang^{a,b}, Hai-peng Liu^{a,b,*}

^a State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen 361102, Fujian, PR China

^b Fujian Collaborative Innovation Center for Exploitation and Utilization of Marine Biological Resources, Fujian Engineering Laboratory of Marine Bioproducts and Technology, Xiamen 361102, Fujian, PR China

ARTICLE INFO

Article history:

Received 25 August 2015

Received in revised form

8 January 2016

Accepted 29 January 2016

Available online 1 February 2016

Keywords:

Crayfish hematopoietic tissue cells

White spot syndrome virus

iTRAQ

Proteomics

ABSTRACT

To elucidate proteomic changes of Hpt cells from red claw crayfish, *Cherax quadricarinatus*, we have carried out isobaric tags for relative and absolute quantitation (iTRAQ) of cellular proteins at both early (1 hpi) and late stage (12 hpi) post white spot syndrome virus (WSSV) infection. Protein database search revealed 594 protein hits by Mascot, in which 17 and 30 proteins were present as differentially expressed proteins at early and late viral infection, respectively. Generally, these differentially expressed proteins include: 1) the metabolic process related proteins in glycolysis and gluconeogenesis, DNA replication, nucleotide/amino acid/fatty acid metabolism and protein biosynthesis; 2) the signal transduction related proteins like small GTPases, G-protein-alpha stimulatory subunit, proteins bearing PDZ- or 14-3-3-domains that help holding together and organize signaling complexes, casein kinase I and proteins of the MAP-kinase signal transduction pathway; 3) the immune defense related proteins such as α -2 macroglobulin, transglutaminase and *trans*-activation response RNA-binding protein 1. Taken together, these protein information shed new light on the host cellular response against WSSV infection in a crustacean cell culture.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

White spot syndrome virus (WSSV) is an enveloped DNA virus with a wide host range in crustaceans such as shrimp, crab and crayfish [1,2]. In shrimp aquaculture WSSV infection will result in collapse of the whole shrimp stock within 3–10 days [3]. After the emergence of the WSSV from 1990s, researchers have been focusing on the virus and the immune response of the host. For example, quantitative determination of the mRNA pool in infected cells or animals versus controls has been widely used [4,5]. With the virus characterized and some knowledge regarding the immune response of the host accumulated [6], a comprehensive approach to investigate how the host physiological system responds to the viral invasion is the important next step to develop efficient control of

the infection. Proteomics is a means of comprehensive interpretation used to describe more direct molecular responses than conventional studies assessing the messenger RNA [7]. Hence, researchers employed proteomic approaches such as two-dimensional electrophoresis for studying host protein changes during WSSV proliferation [8,9]. Recently, iTRAQ combined with liquid chromatography-tandem mass spectrometry is used to directly quantify and compare the protein expression levels of samples with great efficiency and accuracy [10]. Such a global approach helps to correlate different host proteins participating in biological processes, cellular components and molecular function in response to a pathogenic infection.

As one of the principle targets by WSSV, crayfish Hpt cells have been set up as a good model for investigation of WSSV infection [11]. Intriguingly, the crayfish Hpt cells were lately proved by Wu et al. [12] to be a better cell model for WSSV replication than hemocytes which could not afford the virus for proper replication and viral assemble. Previously, we have investigated the transcript alteration of Hpt cells from red claw crayfish, *Cherax quadricarinatus*, towards WSSV challenge using subtractive library

* Corresponding author. State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen 361102, Fujian, PR China.

E-mail address: haipengliu@xmu.edu.cn (H.-p. Liu).

¹ Contributed equally to this work.

screening. We noticed that numerous genes, with differential expression between WSSV infection and mocked infection, were present with annotated functions in immune defense, cytoskeletal organization, signal transduction, stress, metabolism and homeostasis [13]. To further identify proteins or pathways altered during viral infection, here we report proteomic responses of crayfish Hpt cells by iTRAQ at both early (1 hpi) and late (12 hpi) stages post WSSV infection accordingly. These differentially expressed proteins provide us with better understanding of the host's coordinated effort to defend the WSSV infection.

2. Materials and methods

2.1. Viral challenge of crayfish Hpt cells

Adult healthy red claw crayfish were collected from crayfish farms in Zhangpu, Fujian, China. The crayfish were acclimatized for one week at least in well-aerated tanks and only intermolted animals were sacrificed for Hpt cells preparation as described by Söderhäll et al. [14]. The isolated Hpt cells were cultured in six well plates with 85–90% confluence using L-15 medium containing 5 μ M 2-mercaptoethanol, 1 μ M phenylthiourea, 2 mM L-glutamine and a crude astakine preparation [15]. The cells were then inoculated, 1 hpi as early infection stage and 12 hpi as late infection stage accordingly, with 10^5 copies of WSSV (Chinese isolate, WSSV-CNAF332093) kindly provided by Prof. Xun Xu from the Third Institute of Oceanography, State Oceanic Administration, China. UV-inactivated WSSV was used as mocked infection.

2.2. Protein extraction

The cells in duplicates above were harvested by scraping in culture medium with a rubber policeman and collected by centrifugation at 4200 rpm for 5 min at 4 °C. The cell pellets were then lysed with RIPA lysis buffer (Thermo Scientific) and the supernatant was collected after centrifugation followed by protein purification using a 2-D clean-up kit (Bio-Rad), and protein concentration was determined by Lowry method using the RC DC protein assay kit (Bio-Rad).

2.3. iTRAQ labeling and mass spectrometry

For iTRAQ assays, two independent samples were pooled and each pool was divided into duplicates. Total protein (100 μ g) of each sample solution as prepared above was digested with Trypsin Gold (Promega) at a ratio of protein:trypsin = 30:1 at 37 °C for 16 h. Thereafter, peptides were dried by vacuum centrifugation and reconstituted in 0.5 M triethyl ammonium bicarbonate and processed according to the manufacturer's protocol for 8-plex iTRAQ reagent (Applied Biosystems). Two technical replicates were prepared for iTRAQ labeling as suggested by the manufactory instructions. The proteins from the infected (1 hpi labeled with 114/118; 12 hpi labeled with 116/121) and mock-infected samples (1 hpi labeled with 113/117; 12 hpi labeled with 115/119) were labeled with iTRAQ tags (Fig. S1). The tags employed N-hydroxysuccinimide chemistry which consists of three functional groups: an amine-reactive group and an isotopic reporter group (N-methylpiperazine) linked by an isotopic balancer group (carbonyl) for the normalization of the total mass of the tags. The labeled peptide mixtures were pooled and dried by vacuum centrifugation and then reconstituted in buffer A (25 mM NaH₂PO₄ in 25% acetonitrile, pH 2.7) followed by separation on a strong cation exchange chromatography column (4.6 \times 250 mm, 5 μ m, Ultremex SCX, Phenomenex, USA). The peptides were next treated with an isocratic elution at a flow rate of 1 mL/min with buffer A for 10 min followed

by a gradient from 5 to 60% of buffer B (25 mM NaH₂PO₄, 1 M KCl in 25% acetonitrile, pH 2.7) for 27 min, and then a gradient elution from 60 to 100% of buffer B for 1 min. The system was then maintained at 100% buffer B for 1 min before equilibrating with buffer A for 10 min prior to the next injection. Fractions were collected every 1 min and the eluted peptides were pooled into 20 fractions followed by desalting on a Strata X C18 column (Phenomenex) and vacuum-dried.

Each fraction was resuspended in buffer A (5% acetonitrile, 0.1% formic acid) and centrifuged at 20,000 g for 10 min, and the final concentration of peptide was about 0.5 g/L on average. Ten microlitre of supernatant was loaded on a LC-20AD nanoHPLC (Shimadzu, Japan) by the autosampler onto a 2 cm C18 trap column. Then, the peptides were eluted onto a 10 cm analytical C18 column packed in-house. The samples were loaded at 8 μ L/min for 4 min, then the 35 min gradient was run at 300 nL/min starting from 2 to 35% buffer D (95% acetonitrile, 0.1% formic acid), followed by 5 min linear gradient to 60% and by 2 min linear gradient to 80%. After that, the samples were maintained at 80% buffer D for 4 min, and finally return to 5% in 1 min. Data acquisition was performed with a Triple TOF 5600 system (AB SCIEX) fitted with a nanospray III source and a pulled quartz tip as the emitter (New Objectives, MA). Data was acquired using an ion spray voltage of 2.5 kV, curtain gas of 30 psi, nebulizer gas of 15 psi, and an interface heater temperature of 150 °C. The MS was operated with a RP of greater than or equal to 30,000 FWHM for TOF MS scan. For IDA, survey scans were acquired in 250 ms and as many as 30 product ion scans were collected if they exceeded a threshold of 120 counts per second. Total cycle time was fixed at 3.3 s. The Q2 transmission window was 100 Da for 100%. Four time bins were summed for each scan at a pulser frequency value of 11 kHz through monitoring of the 40 GHz multichannel TDC detector with four anode channel detect ion. A sweeping collision energy setting of 35 \pm 5 eV, coupled with iTRAQ adjust rolling collision energy, was applied to all precursor ions for collision induced dissociation. Dynamic exclusion was set for half of peak width (15 s) and then the precursor was refreshed off the exclusion list.

2.4. Database search and bioinformatic analysis

Peptide sequence data were processed using Mascot software version 2.3.02 to obtain the information of associated proteins. A false discovery rate of less than 1% was used as cut off. The proteins were further classified using Gene Ontology (GO) and pathway enrichment analysis (<http://www.geneontology.org>). To determine the alteration of protein expression induced by WSSV infection, protein expression fold changes were calculated relatively in WSSV infected Hpt cells compared to those of mock-infected cells. Student's t-test was used to identify statistically significant changes in protein expression ($p < 0.05$). The quantitative proteins ratios of ≥ 1.3 were considered as up-regulation and proteins with a label ratio of ≤ 0.77 [16,17] were considered as down-regulated proteins in WSSV infected Hpt cells versus mock-infected cells.

2.5. Quantitative real-time polymerase chain reaction

RNA isolation was performed by GenElute mammalian total RNA isolation kit (Sigma-Aldrich) according to manufacturers protocol. Briefly, reverse transcription was done with 1 μ g of total RNA using Superscript II Reverse Transcriptase (Invitrogen) according to instructions of manufacturer. Real-time PCR was performed on ABI 7500 using FastStart Universal SYBR master (Roche) and the primers used were included in supplementary information (Table S2). The expression level of mRNA was normalized by 16S rRNA. The data was presented as relative expression levels (means \pm SD,

$n = 3$), and subjected to one-way Analysis of Variance (ANOVA) followed by Fisher's least significant difference test. The significant difference between WSSV treated and control (UV treated WSSV) samples was represented by one asterisk for $p < 0.05$.

3. Results and discussion

3.1. Protein profiling and iTRAQ quantification

To get insight into the proteins differentially expressed at both early stage (1 hpi) and late stage (12 hpi) of WSSV infection as recent publications reported that these two time points have been clearly defined accordingly of WSSV replication in crayfish hpt cells [12,13], iTRAQ analysis was employed for quantification followed by protein profiling, which resulted in 594 protein hits in Mascot. The functional enrichment analysis of GO annotations for biological processes showed that the identified proteins were classified into 23 categories (Fig. 1), in which proteins related to cellular process (17.14%) and metabolic process (15.41%) were the most abundant molecules followed by cellular component organization biogenesis (6.43%), biological regulation (6.37%), regulation of biological process (5.85%), developmental process (5.44%), response to stimulus (4.90%), signaling transduction (2.95%), negative regulation of biological process (2.23%) and others. Apparently, the protein expression in Hpt cells showed variation at both early and late stages of infection if compared to those of mock-infected cells (Tables 1–3). Among these proteins, alpha amino adipic semialdehyde synthase, ribonucleoside-diphosphate reductase and thymidylate synthase in metabolism; casein kinase I isoform alpha, cAMP-dependent protein kinase, PDZ domain containing protein GIPC1 and Rab-1 involved in signal transduction were found for the first time to be involved in the host response to WSSV infection. Furthermore, the cytoskeletal associated proteins like calponin-3 and beta chain spectrin were also found to be significantly expressed towards WSSV infection. Here we focus on the brief interpretation of diverse functional classes related to metabolic processes, signal transduction proteins and immune related proteins, which are putatively key molecules involved in metabolism and host defense against cellular stress caused by WSSV infection. In addition, numerous proteins were expressed in WSSV infected cells but without significant difference in up- or down-regulation when compared to a mock infection (Table S1).

3.2. Metabolic process related proteins

Viruses require host cellular metabolism for their energy needs and successful replication. WSSV infection perturbs the expression

of at least 14 enzymes action in glycolysis, gluconeogenesis, the pentose phosphate pathway, metabolism of nucleotide, amino acid and fatty acid, DNA replication and protein biosynthesis. An increase in activity of metabolic pathways associated with glycolysis, the pentose phosphate pathway, nucleotide biosynthesis, glutaminolysis and amino acid biosynthesis were observed in shrimp hemocytes post WSSV challenge [18]. In our present study, phosphoglycerate mutase 2 with a role in glycolysis was down-regulated at both 1 hpi and 12 hpi post infection. Further studies with glucose starved media should be done to ascertain its role in WSSV replication and possible therapeutic applications. The pentose phosphate pathway fulfills two essential functions, the formation of pentose phosphate for synthesis of nucleotides, RNA, DNA and the generation of NADPH for biosynthetic reactions to maintain glutathione in the reduced state and protecting the cell from damaging reactive oxygen intermediates. Glucose-6-phosphate dehydrogenase, the key enzyme of pentose phosphate pathway showed increased activity in WSSV infected *Litopenaeus vannamei* [19]. Interestingly, in our study the key enzyme of the oxidative pentose phosphate pathway, 6-phosphogluconate dehydrogenase, was up-regulated at 12 hpi whereas in *Penaeus monodon* it was down-regulated during YHV infection [20]. This difference may reflect alternative energy utilization of these two viruses.

Viruses use several strategies to ensure adequate supply of nucleotide by expressing virus encoded enzymes that boost nucleotide synthesis and increasing the flux of pentose phosphate pathway [21–23]. WSSV-infection perturbs the nucleotide biosynthesis, particularly on nucleoside and nucleotide metabolic enzymes involved in producing monomer intermediates for DNA synthesis such as ribonucleoside-diphosphate reductase, thymidylate synthase and GMP synthase. The first two enzymes were down-regulated at 1 hpi whereas GMP synthase, an ATP-dependent enzyme involved in purine nucleotide synthesis, and thymidylate synthase was over-expressed at 12 hpi (in comparison to its expression at 1 hpi with live virus infection, Table 3). A similar increase in nucleotide biosynthesis was observed during WSSV infection in *L. vannamei* [18]. The relative increase of nucleosides at 12 hpi when compared to 1 hpi implies that the DNA replication machinery might be sequestered to produce viral DNA. Recently pharmaceutical compounds are proved to inhibit RNA, DNA and retroviruses by targeting pyrimidine biosynthesis [24]. Broad-spectrum antivirals targeting nucleotide biosynthesis could be further tested in vitro such as in the Hpt cell cultures. Antiviral drugs targeting nucleotide biosynthesis is commonly used against human viruses and this might be extended to WSSV. Differential expression of enzymes associated with amino acid metabolism was observed in our present study. Alpha amino adipic semialdehyde




Fig. 1. GO analysis of total number of proteins involved in biological processes identified in crayfish Hpt cells post WSSV infection.

Table 1

The significant change in protein expression of Hpt cells post WSSV challenge at 1 hpi with respect to UV-inactivated WSSV.

Protein name	Accession	Score	Coverage (%)	Unique peptide	Fold change
Metabolic process					
α -aminoacidic semialdehyde synthase	B0W052	279	1.9	1	0.6
Aspartate aminotransferase	F5A6C3	121	6	1	1.42
Phosphoglycerate kinase	H9K1M9	223	3.5	1	1.33
Phosphoglycerate mutase	G0ZJ11	163	17	3	1.4
Phosphoglycerate mutase 2	B0X5F6	122	3.5	1	0.5
Ribonucleoside-diphosphate reductase	E9GDU6	106	4.9	2	0.58
Thymidylate synthase	E0VDQ0	70	2.2	1	0.33
Signal transduction					
Mitogen-activated protein kinase kinase	K7XD34	152	9.1	2	0.52
Proteasome subunit alpha type	E9GIX9	186	13.7	2	0.74
Rab-1	L7RUN6	692	46.6	4	1.4
Ras opposite	D6X469	156	4.3	2	0.7
Immune related					
α 2-macroglobulin 2	G0ZJ42	332	33	4	0.7
Glutathione peroxidase	C5H3W2	532	11.4	1	1.38
Transglutaminase	Q964D3	3536	4.2	1	0.7
T-complex protein 1 subunit zeta	B0W8W8	84	2.3	1	1.4
Cytoskeletal					
Beta chain spectrin-like protein	F8QXF5	220	13.9	2	0.68
Calponin-3	C1BUD8	203	6.9	1	0.66

Table 2

The significant change in protein expression of Hpt cells post WSSV challenge at 12 hpi with respect to UV-inactivated WSSV.

Protein name	Accession	Score	Coverage (%)	Unique peptide	Fold change
Metabolic process					
6-phosphogluconate dehydrogenase	E9GAM5	676	6	2	1.6
Enoyl-CoA hydratase	F0J8W9	153	9.4	1	0.68
GF15148	B3MMZ7	82	4.4	1	0.6
GMP synthase	E1ZYK1	205	2.5	1	2.32
Phosphoglycerate mutase 2	B0X5F6	122	3.5	1	0.7
Tudor staphylococcal nuclease	G1ARD5	311	4.5	3	0.6
Signal transduction					
14-3-3 epsilon	Q1HPT4	1248	21	1	1.45
26 S proteasome regulatory subunit rpn2	B7Q0A9	153	1.3	1	0.71
cAMP-dependent protein kinase type II regulatory subunit	E2ACB7	101	2.9	1	0.76
Casein kinase I isoform alpha	E2A2Y7	116	5.9	2	1.46
GTP binding protein alpha subunit Gs	C7G305	124	5.5	1	1.47
Mitogen-activated protein kinase 14	G1UHC5	181	13.4	4	0.7
PDZ domain-containing protein GIPC1	C1C1V8	107	5	1	2.86
Immune related					
α 2-macroglobulin	A0T1M1	74	0.5	1	1.4
Glutathione peroxidase	C5H3W2	532	11.4	1	0.77
Glutathione S-transferase	G0ZJ66	89	6.9	1	1.42
Hemocytin	G0ZJ16	321	25.7	3	0.77
T-complex protein 1 subunit zeta	B0W8W8	84	2.3	1	0.6
Cytoskeletal					
65-kDa macrophage protein	B7QBZ0	187	2.2	1	1.38

Table 3

The significant change in protein expression of Hpt cells post WSSV challenge at 12 hpi with respect to that of expression at 1 hpi.

Protein name	Accession	Score	Coverage (%)	Unique peptide	Fold change
Metabolic process					
α -aminoacidic semialdehyde synthase	B0W052	279	1.9	1	0.6
Adenosylhomocysteinase	A0JCJ9	146	3.5	1	1.47
Bifunctional aminoacyl-tRNA synthetase	G0ZJ24	77	15.4	2	1.3
Thymidylate synthase	E0VDQ0	70	2.2	1	1.56
Signal transduction					
cAMP-dependent protein kinase type II regulatory subunit	E2ACB7	101	2.9	1	0.65
Mitogen-activated protein kinase kinase	K7XD34	152	9.1	2	1.49
Immune related					
α 2-macroglobulin 2	G0ZJ42	332	33	4	1.51
Protein spätzle	G0ZJ18	728	49.7	6	1.49
T-complex protein 1 subunit eta	E2BZD9	283	7.7	1	1.33
Cytoskeletal					
Ca ²⁺ sensor	B7PX38	289	29.8	5	0.72

synthase, involved in the lysine degradation pathway was down-regulated at 1 hpi while increased to control levels at 12 hpi. Adenosylhomocysteinase, a key enzyme in the regulation of the intracellular concentration of S-adenosylhomocysteine (SAH) was up-regulated at 12 hpi (Table 3). SAH is an intermediate in the synthesis of cysteine and adenosine. An increased intracellular concentration of SAH inhibited vaccinia virus replication in murine cells [25]. Aminoacyl-tRNA synthetases (ARSs) are highly conserved for efficient and precise translation of genetic codes. Evidence is accumulating that several ARSs form a macromolecular protein complex that would work as a molecular hub linked to the multiple signaling pathways [26]. The up-regulation of bifunctional aminoacyl-tRNA synthetase at 12 hpi (Table 3) implies that protein synthesis might be elevated during later stages of the WSSV infection. Recent studies in crayfish found that there was a decrease in long chain fatty acids during WSSV infection at 12 hpi and later it increased at 24 hpi [27]. The knowledge of virus induced changes in cell metabolome may lead to potential use of inhibitors of metabolic enzymes to treat WSSV infection.

3.3. Response of signal transduction related proteins

In total eleven proteins related to signal transduction were differentially expressed in Hpt cells upon WSSV-infection. Three of those are GTP-binding proteins (Rab-1, Ras opposite, GTP-alpha G_s). GTP binding proteins regulate a wide variety of cell functions acting as biological timers, for instance, that initiate and terminate specific cell functions. In addition they play key roles in determining spatial locations of specific cell functions. Rab-1 regulates protein transport from endoplasmic reticulum to Golgi apparatus. Rab GTPase is essential for the control of intracellular membrane trafficking in all eukaryotic cells and further affects the ability of phagocytic cells to scavenge pathogen. This is in synchrony with the observed up-regulation of Rab-1 protein during 1 hpi of the current study. In *Penaeus monodon*, Rab 7 was found to bind to rVP28 of WSSV. The in vivo neutralization assay of WSSV infected shrimp in the presence of anti-Rab7 antibody revealed less mortality compared to that of WSSV alone [28]. Depletion of Rab-1 significantly decreased the replication of Hepatitis C virus in a human hepatoma cell line [29]. The upregulation of Rab-1 suggests its role in WSSV pathogenesis. G-protein alpha stimulatory subunit (G_{α_s}) is activated by G-protein coupled receptors located in the cell membrane. G_{α_s} stimulates adenylate cyclase to produce cAMP [30] a ubiquitous second messenger that enables cells to detect and respond to extracellular signals [31]. The intracellular cAMP-level affects the cAMP-dependent protein kinase (PKA) and consequently many metabolic processes of the cell. In our present study WSSV-infection caused over expression of G_{α_s} at 12 hpi. More in depth investigation is needed to understand its role during a late infection stage of WSSV infection.

Adapter molecules or domains such as GIPC1 and 14-3-3 are involved in protein-protein interaction. In the current study the strongest response was the expression of GIPC1, a scaffolding protein that regulates cell surface receptor expression and trafficking. Viruses usually use carboxyl-terminal PDZ domain-binding motif (PBM) that mediates interactions with cellular PDZ proteins. GIPC1 contains such a PDZ domain, a protein-protein interaction module typically found in cytoplasmic and membrane adapter proteins that are involved in a variety of cellular processes of significance to viral infection, for instance maintenance of cell-cell junctions, cellular polarity, and signal transduction pathways [32]. GIPC1 was in control level at first hour after viral challenge and later up-regulated at 12 hpi, implying that the GIPC1 might have modulated the cellular process in favor of WSSV replication and dissemination in the cells. To reveal the role of GIPC1 during WSSV

infection, protein-protein interaction studies should be undertaken in the future. 14-3-3 epsilon protein is known for its ability to bind multiple cellular protein ligands. It interacts with a wide range of proteins for functions like cell cycle control, regulation of cell death and apoptosis etc. In addition 14-3-3 epsilon can interact with a diverse array of viral products, and thereby re-direct cellular processes [33–35]. For example, Hepatitis C virus core protein can interact with 14-3-3 epsilon protein [33]. Also severe acute respiratory syndrome corona virus nucleocapsid protein was phosphorylated and localized in the cytoplasm by 14-3-3-mediated translocation [36]. In our study 14-3-3 epsilon was at control levels at 1 hpi and up-regulated at 12 hpi and therefore indicative of an established viral infection. Further studies are therefore needed to elucidate the interaction of 14-3-3 with viral proteins.

There was a trend of down regulation observed in proteins that assemble the proteasome (Proteasome subunit alpha type, 26S proteasome regulatory subunit rpn2). The main function of proteasome is to degrade unneeded or damaged proteins. There is a high stoichiometric imbalance of host target proteins over those encoded by the virus. To overcome this, many viruses have evolved mechanisms by which their cellular target proteins are directed to the 26S proteasome and subjected to proteolytic degradation [37]. Recently a plethora of viral proteins have been shown to direct host-cell proteins for proteolytic degradation. These activities are required for various aspects of viral life cycle like viral entry [38], replication [39], enhanced survival [40,41], and to viral release [42]. Similar to our study, infection of enterovirus 61 in rhabdomyosarcoma cells resulted in down-regulation of proteasome subunit alpha type 2 [43].

The cellular signal transduction is largely controlled by protein phosphorylation. Several protein kinases were differentially expressed during WSSV infection such as protein casein kinase I (CK1) isoform alpha, a serine/threonine kinase, which was up-regulated at 12 hpi. It inhibits hyperphosphorylation of nonstructural protein 5A of hepatitis C virus. The nonstructural (NS) protein 5A is a phosphoprotein and experiments indicated that the phosphorylation state of NS5A is important for the outcome of viral RNA replication [44]. Similarly, in our study the up-regulation of CK1 might be indicative of its interaction with WSSV proteins. WSSV envelope proteins like VP28 and VP19 were previously reported to be threonine phosphorylated [45]. Further investigations are needed to understand the role of casein kinase I isoform alpha. cAMP-dependent protein kinase (PKA) is involved in the regulation of glycogen, sugars and lipid metabolism. Its down-regulation at 12 hpi (Table 3) might affect the production of essential metabolites during infection. Activation of cAMP can block apoptosis induced by Herpes simplex virus 1 associated genes [46]. As the WSSV infection is nearly established at 12 hpi, WSSV might suppress cAMP activation for promoting apoptosis in infected cells, spreading viral infection to nearby cells. The mitogen-activated protein kinase (MAPK) cascade pathways have global role in the intracellular signaling network pathway. Mitogen-activated protein kinase (MAP2K/MEK) phosphorylates mitogen-activated protein kinase (MAPK/ERK) are part of the protein kinase cascade starting with an extracellular stimulant ending with the expression of transcription factors in the nucleus. It is thought that a viral infection acts as an extracellular stimulant that activates this pathway. RNA silencing of p38 MAPK homologue *ivp38* caused increase in WSSV replication on *L. vannamei* [47]. In our study MAPK was in control levels at 1 hpi but down-regulated at 12 hpi while MAP2K down-regulated at initial period and up-regulated at latter time point (Table 3). The significant expression of MAPK and MAP2K should be further investigated.

3.4. Response of immune related proteins

Among these proteins, nine immune related proteins were differentially expressed during this period of infection within 12 hpi. Two of these proteins, α 2-macroglobulin and transglutaminase, are well known to play a role in coagulation of hemolymph during pathogen infection. For instance, α 2-macroglobulin is considered as a broad range proteinase inhibitor involved in phagocytosis [48,49]. Silencing of α 2-macroglobulin clearly reduced phagocytosis of *Chryseobacterium indologenes* by hemocytes both in vitro and in vivo in hard tick *Ixodes ricinus* [50]. According to the iTRAQ quantification, no change in protein expression was found for α 2-macroglobulin at 1 hpi but its expression was clearly up-regulated at 12 hpi, suggesting that α 2-macroglobulin might have a role during late infection hours of WSSV in crayfish Hpt cells. Transglutaminase has a regulatory function in antimicrobial peptide production in kuruma shrimp. Silencing of transglutaminase caused the down regulation of antimicrobial peptides such as crustin and lysozyme, thus, negatively affecting on the host's defense against pathogen infection [51]. Additionally, it has been documented that in a freshwater crayfish transglutaminase prevents hematopoietic stem cells from migrating into the hemolymph [52]. As hemocytes are particularly important for innate immune defense in invertebrates, this prevention on release of matured Hpt cells from the hematopoietic tissue is likely to have an impact on host immunity. In our present study transglutaminase expression was down-regulated at 1 hpi if compared to those of mock-infection, implying that transglutaminase might act a critical role during virus infection. Further functional studies on transglutaminase against WSSV infection in red claw crayfish both in vitro and in vivo will benefit the elucidation of an important role of this enzyme.

Späetzle is the well-known component of Toll signaling pathway, which plays a key role in regulating innate immune responses in invertebrates. Gram positive bacteria, fungi and certain viruses can activate Toll signaling pathway initiating an intracellular signaling cascade to promote the expression of immune-related genes such as antimicrobial peptides through the activation of the NF- κ B family-Dorsal proteins [53]. It has been shown that key components of the Toll pathway, like Toll [54] and dorsal [55], are activated during viral infection in shrimp. By recognition of high mannose residues expressed on the surface of viruses or virus surface glycoproteins, Toll like receptors are able to mediate virus entry into host cells, in which binding of ligand späetzle to Toll is necessary to initiate this pathway [56]. Recently, transcriptomic changes of späetzle was also reported during WSSV and *Vibrio* infection in *Fenneropenaeus chinensis* [57]. Protein späetzle was up-regulated in crayfish Hpt cells at 12 h post WSSV infection (Table 3). Recent studies in crayfish Hpt shows that by 12 hpi progeny WSSV virions will be successfully generated [12]. The presence of progeny virions at this stage might be the reason for up regulation of protein späetzle. Further functional studies are needed to understand the late expression of späetzle during WSSV infection.

Antioxidant enzymes regulate the concentrations of radical oxygen species, for instance, glutathione peroxidase detoxifies lipids and hydrogen peroxidase while glutathione-S-transferase marks xenobiotics for cellular degradation. This antioxidant activity usually is increased upon pathogenic infection. We found that glutathione peroxidase and glutathione-S-transferase were obviously affected by WSSV infection in crayfish Hpt cells, in which the former expression was promoted at 1 hpi and the latter was enhanced at 12 hpi. An increased glutathione peroxidase activity at early hours and an up-regulated glutathione-S-transferase activity at late hours were also found in *L. vannamei* post WSSV infection [58]. Increased oxidative damage will occur in response to the

challenge when rate of free radical formation exceeds its removal by antioxidant system [58]. Faulty antioxidant defense causes the failure of sensitive cellular and subcellular components leading to cell death [59]. These findings clearly indicated that antioxidant response was modulated during WSSV infection in crustaceans [58].

Invertebrates and plants are capable of suppressing viral infection through RNA silencing mediated by RISC. As a key component of RISC, *trans*-activation response RNA-binding protein (TRBP) was found to be associated with Dicer and Argonautes, both of which are involved in RNA interference and microRNA pathways [60]. Combination of TRBP with eukaryotic initiation factor 6 leads to an antiviral RNA interference pathway in shrimp *Marsupenaeus japonicus* [61]. Knockdown of a shrimp tudor staphylococcal nuclease (PmTSN), one of the RISC associated protein involved in post transcriptional regulation, resulted in reduced inhibition of YHV infection mediated by RNAi of YHV in *Penaeus monodon* [62]. In our studies we found that TAR RNA-binding protein isoform 1 and TSN, both involved in RISC, were down-regulated at 12 hpi. This implied that WSSV-infection might negatively affect the immune defense against this viral infection via RNAi process, but how this effect is elicited and modulated during WSSV replication needs further exploration.

3.5. Validation of selected proteins by quantitative real-time polymerase chain reaction

To validate the differentially expressed proteins identified by iTRAQ, a randomly selected group of proteins transcriptional level was confirmed by quantitative real-time PCR. As shown in Fig. 2, the gene expression levels were in agreement to our iTRAQ results. Among these tested genes, tudor staphylococcal nuclease and thymidylate synthase involved in metabolic process group showed similar gene expression trend to our proteomics data. Briefly, tudor staphylococcal nuclease showed a significant decrease at 12 hpi, while thymidylate synthase was significantly down-regulated at 1 hpi but up-regulated at 12 hpi. In regarding to the signal transduction group, the transcriptional level of MAPKK, MAPK-14, 14-3-3 epsilon and ras opposite were analyzed. MAPKK was significantly decreased at 1 hpi but increased at late infection stage, and MAPK-14 was clearly down-regulated at late hours of infection. Ras opposite showed a decrease on initial hours while 14-3-3 epsilon was obviously enhanced at late hours of infection. In case of immune-related and cytoskeletal group, both transglutaminase and calponin-3 showed significant decline at initial hours post WSSV infection, which were in well consistent with our iTRAQ result.

4. Conclusion

The WSSV infection in the crayfish Hpt cells obviously altered the host cellular expression of proteins related to metabolic processes, signal transduction, immunity and cytoskeletal organization. WSSV utilized host metabolic system for its energy needs, replication and dissemination inside the host cells. The virus managed to alter important signal transduction proteins those function as GTP binding, adapter molecules, protein biosynthesis and protein phosphorylation. The protein expression in early hours exposes vulnerability of host defense mechanism to the viral infection. The role of many of the identified proteins in viral infection remains unknown, some of those are first observations and will require systematic studies to show the mechanism of action. By identifying a series of proteins that are affected by viral infection this study opens new opportunities in two directions: 1) to better understand host-virus interactions and 2) to select targets




Fig. 2. Validation of proteomic data using RT-qPCR. Relative mRNA levels (y axis) of WSSV treated hpt cells at 1 hpi and 12 hpi where UV treated WSSV was used as control. Selected genes validated were: a) Tudor staphylococcal nuclease; b) Thymidylate synthase; c) MAPKK; d) MAPK-14; e) 14-3-3 epsilon; f) Ras opposite; g) Transglutaminase; h) Calponin - 3.

to reduce vulnerability of the host toward the challenge. The proteins identified and host physiological changes elucidated can be further utilized for developing putative drug targets for controlling this virus.

Acknowledgments

This study was supported by an 863 program (2012AA092205), NSFC (31222056, 41176114, 41476117, J1210050), NCET-10-0711, FRFCU (20720150155) and NSFFJ (2013J06010).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.fsi.2016.01.035>.

References

- [1] S.C. Chen, D.H. Jones, E. Fynan, G.H. Farrar, J.C. Clegg, H.B. Greenberg, et al., Protective immunity induced by oral immunization with a rotavirus DNA vaccine encapsulated in microparticles, *J. Virol.* 72 (1997) 5757–5761.
- [2] C.F. Lo, C.H. Ho, S.E. Peng, C.H. Chen, H.C. Hsu, Y.L. Chiu, et al., White spot syndrome baculovirus (WSBV) detected in cultured and captured shrimp, crabs and other arthropods, *Dis. Aquat. Organ.* 27 (1996) 215–226.
- [3] D.V. Lightner, *A Handbook of Shrimp Pathology and Diagnostic Procedures for Disease of Cultured Penaeid Shrimp*, World Aquaculture Society, Baton Rouge, LA, 1996.
- [4] A. Tassanakajon, K. Somboonwivat, P. Supungul, S. Tang, Discovery of immune molecules and their crucial functions in shrimp immunity, *Fish. Shellfish Immunol.* 34 (2013) 954–967.
- [5] T. Flegel, K. Sritunyalucksana, Shrimp molecular responses to viral pathogens, *Mar. Biotechnol.* 13 (2011) 587–607.
- [6] H.P. Liu, K. Söderhäll, P. Jiravanichpaisal, Antiviral immunity in crustaceans, *Fish. Shellfish Immunol.* 27 (2009) 79–88.
- [7] N.L. Anderson, N.G. Anderson, Proteome and proteomics: new technologies, new concepts, and new words, *Electrophoresis* 19 (1988) 1853e61.
- [8] Y.M. Chai, S.S. Yu, X.F. Zhao, Q. Zhu, J.X. Wang, Comparative proteomic profiles of the hepatopancreas in *Fenneropenaeus chinensis* response to white spot syndrome virus, *Fish. Shellfish Immunol.* 29 (2010) 480–486.
- [9] A.D. Kulkarni, V. Kiron, J.H.W.M. Rombout, M.F. Brinchmann, J.M.O. Fernandes, N.S. Sudheer, et al., Protein profiling in the gut of *Penaeus monodon* gavaged with oral WSSV-vaccines and live white spot syndrome virus, *Proteomics* 14 (2014) 1660–1673.
- [10] C. Evans, J. Noirel, S.Y. Ow, M. Salim, A.G. Pereira-Medrano, N. Couto, et al., An insight into iTRAQ: where do we stand now? *Anal. Bioanal. Chem.* 404 (2012) 1011–1027.
- [11] P. Jiravanichpaisal, K. Söderhäll, I. Söderhäll, Characterization of white spot syndrome virus replication in *in vitro*-cultured haematopoietic stem cells of freshwater crayfish, *Pacifastacus leniusculus*, *J. Gen. Virol.* 87 (2006) 847–854.
- [12] J. Wu, F. Li, J. Huang, L. Xu, F. Yang, Crayfish hematopoietic tissue cells but not hemocytes are permissive for white spot syndrome virus replication, *Fish. Shellfish Immunol.* 43 (2015) 67–74.
- [13] H.P. Liu, R.Y. Chen, Q.X. Zhang, H. Peng, K.J. Wang, Differential gene expression profile from haematopoietic tissue stem cells of red claw crayfish, *Cherax quadricarinatus*, in response to WSSV infection, *Dev. Comp. Immunol.* 35 (2011) 716–724.
- [14] I. Söderhäll, E. Bangyeekhun, S. Mayo, K. Söderhäll, Hemocyte production and maturation in an invertebrate animal; proliferation and gene expression in hematopoietic stem cells of *Pacifastacus leniusculus*, *Dev. Comp. Immunol.* 8 (2003) 661–672.
- [15] I. Söderhäll, Y.A. Kim, P. Jiravanichpaisal, S.Y. Lee, K. Söderhäll, An ancient role for a prokineticin domain in invertebrate hematopoiesis, *J. Immunol.* 174 (2005) 6153–6160.
- [16] R.D. Unwin, J.R. Griffiths, A.D. Whetton, Simultaneous analysis of relative protein expression levels across multiple samples using iTRAQ isobaric tags with 2D nano LC-MS/MS, *Nat. Protoc.* 5 (2010) 1574–1582.
- [17] C.M. Warren, D.L. Geenen, D.L.J. Helseth, H. Xu, R.J. Solaro, Sub-proteomic fractionation, iTRAQ, and OFFGEL-LC-MS/MS approaches to cardiac proteomics, *J. Proteom.* 73 (2010) 1551–1561.
- [18] M.A. Su, Y.T. Huang, I.T. Chen, D.Y. Lee, Y.C. Hsieh, C.Y. Li, et al., An invertebrate Warburg effect: a Shrimp virus achieves successful replication by altering the host Metabolome via the PI3K-Akt-mTOR pathway, *PLoS Pathog.* 10 (2014) e1004196.
- [19] I.T. Chen, T. Aoki, Y.T. Huang, I. Hirono, T.C. Chen, J.Y. Huang, et al., White spot syndrome virus induces metabolic changes resembling the warburg effect in shrimp hemocytes in the early stage of infection, *J. Virol.* 85 (2011) 12919–12928.
- [20] A. Bourchookarn, P.O. Havanapan, V. Thongboonkerd, C. Krittanai, Proteomic analysis of altered proteins in lymphoid organ of yellow head virus infected *Penaeus monodon*, *Biochim. Biophys. Acta* 1784 (2008) 504–511.
- [21] J. Munger, B.D. Bennett, A. Parikh, X.J. Feng, J. McArdle, H.A. Rabitz, et al., Systems-level metabolic flux profiling identifies fatty acid synthesis as a target for antiviral therapy, *Nat. Biotechnol.* 26 (2008) 1179–1186.
- [22] J.B. Ritter, A.S. Wahl, S. Freund, Y. Genzel, U. Reichl, Metabolic effects of influenza virus infection in cultured animal cells: intra- and extracellular metabolite profiling, *BMC Syst. Biol.* 4 (2010) 61.
- [23] T. Daikoku, N. Yamamoto, K. Maeno, Y. Nishiyama, Role of viral ribonucleotide reductase in the increase of dTTP pool size in herpes simplex virus-infected Vero cells, *J. Gen. Virol.* 72 (1991) 1441–1444.
- [24] H.H. Hoffmann, A. Andrea Kunza, V.A. Simona, P. Palesea, M.L. Shawa, Broad-spectrum antiviral that interferes with de novo pyrimidine biosynthesis, *Proc. Natl. Acad. Sci. U. S. A.* 108 (2011) 5777–5782.
- [25] M. Hasobe, J.M. Mckee, R.T. Borchardt, Relationship between intracellular concentration of S-adenosylhomocysteine and inhibition of vaccinia virus replication and inhibition of murine L-929 cell growth, *Antimicrob. Agents Chemother.* 33 (1989) 828–834.
- [26] S.G. Park, E.C. Choi, S. Kim, Aminoacyl-tRNA synthetase-interacting multi-functional proteins (AIMPs): a triad for cellular homeostasis, *IUBMB Life* 62 (2010) 296–302.
- [27] Y.C. Hsieh, Y.M. Chen, C.Y. Li, Y.H. Chang, S.Y. Liang, S.Y. Lin, et al., To complete its replication cycle, a shrimp virus changes the population of long chain fatty acids during infection via Q4 the PI3K-Akt-mTOR-HIF1 α pathway, *Dev. Comp. Immunol.* 53 (2015) 85–95.
- [28] K. Sritunyalucksana, W. Wannapapho, C.F. Lo, T.W. Flegel, PmRab7 is a VP28-binding protein involved in white spot syndrome virus infection in shrimp, *J. Virol.* 80 (2006) 10734–10742.
- [29] E.H. Sklan, R.L. Serrano, S. Einav, S.R. Pfeffer, D.G. Lambright, J.S. Glenn, TBC1D20 is a Rab1 GTPase-activating protein that mediates hepatitis C virus replication, *J. Biol. Chem.* 282 (2007) 36354–36361.
- [30] J.J. Tesmer, R.K. Sunahara, A.G. Gilman, S.R. Sprang, Crystal structure of the catalytic domains of adenylyl cyclase in a complex with Gs, *Science* 278 (1997) 1907–1916.
- [31] J.A. Alspaugh, R. Pukkila-Worley, T. Harashima, L.M. Cavallo, D. Funnell, G.M. Cox, et al., Adenylyl cyclase functions downstream of the G α protein Gpa1 and controls mating and pathogenicity of *Cryptococcus neoformans*, *Eukaryot. Cell* 1 (2002) 75–84.
- [32] R.T. Javier, A.P. Rice, Emerging theme: cellular PDZ proteins as common targets of pathogenic viruses, *J. Virol.* 85 (2011) 11544–11556.
- [33] H. Aoki, J. Hayashi, M. Moriyama, Y. Arakawa, O. Hino, Hepatitis C virus core protein interacts with 14-3-3 protein and activates the kinase Raf-1, *J. Virol. Methods* 74 (2000) 1736–1741.
- [34] S.I. Han, M.A. Kawano, K. Ishizu, H. Watanabe, M. Hasegawa, S.N. Kanesashi, Rep68 protein of adeno-associated virus type 2 interacts with 14-3-3 proteins depending on phosphorylation at serine 535, *Virology* 320 (2004) 144–155.
- [35] Z. Pei, M.S. Harrison, A.P. Schmitt, Parainfluenza virus 5 m protein interaction with host protein 14-3-3 negatively affects virus particle formation, *J. Virol.* 85 (2011) 2050–2059.
- [36] M. Surjit, R. Kumar, R.N. Mishra, M.K. Reddy, V.T. Chow, S.K. Lal, The severe acute respiratory syndrome coronavirus nucleocapsid protein is phosphorylated and localizes in the cytoplasm by 14-3-3-mediated translocation, *J. Virol.* 79 (2005) 11476–11486.
- [37] L. Banks, D. Pim, M. Thomas, Viruses and the 26S proteasome: hacking into destruction, *Trends Biochem. Sci.* 28 (2003) 452–459.
- [38] R. Galinier, E. Gout, H. Lortat-Jacob, J. Wood, J. Chroboczek, Adenovirus protein involved in virus internalisation recruits ubiquitin-protein ligases, *Biochemistry* 41 (2002) 14299–14305.
- [39] J. Parkinson, S.P. Lees-Miller, R.D. Everett, Herpes simplex virus type 1 immediate-early protein vmw110 induces the proteasome-dependent degradation of the catalytic subunit of DNA-dependent protein kinase, *J. Virol.* 73 (1999) 650–657.
- [40] M. Thomas, D. Pim, L. Banks, The role of the E6-p53 interaction in the molecular pathogenesis of HPV, *Oncogene* 18 (1999) 7690–7700.
- [41] L. Coscoy, D.J. Sanchez, D. Ganem, A novel class of herpesvirus-encoded membrane-bound E3 ubiquitin ligases regulates endocytosis of proteins involved in immune recognition, *J. Cell Biol.* 155 (2001) 1265–1273.
- [42] R.N. Harty, M.E. Brown, J.P. McGettigan, G. Wang, H.R. Jayakar, J.M. Huijbregtse, et al., Rhabdoviruses and the cellular ubiquitin-proteasome system: a budding interaction, *J. Virol.* 75 (2001) 10623–10629.
- [43] W.F. Leong, V.T.K. Chow, Transcriptomic and proteomic analyses of rhabdomyosarcoma cells reveal differential cellular gene expression in response to reovirus 71 infection, *Cell. Microbiol.* 8 (2006) 565–580.
- [44] M. Quintavalle, S. Sambucini, V. Summa, L. Orsatti, F. Talamo, R. De Francesco, Hepatitis C virus NS5A is a direct substrate of casein kinase I- α , a cellular kinase identified by inhibitor affinity chromatography using specific NS5A hyperphosphorylation inhibitors, *J. Biol. Chem.* 282 (2007) 5536–5544.
- [45] X. Xie, L. Xu, F. Yang, Proteomic analysis of the major envelope and nucleocapsid proteins of white spot syndrome virus, *J. Virol.* 80 (2006) 10615–10623.
- [46] L. Benetti, B. Roizman, Herpes simplex virus protein kinase US3 activates and functionally overlaps protein kinase A to block apoptosis, *PNAS* 101 (2004) 9411–9416.
- [47] H. Shi, L. Ruan, X. Yan, D. Yao, X. Xu, The role of *Litopenaeus vannamei* p38 in white spot syndrome virus infection, *Dev. Comp. Immunol.* 44 (2014) 180–185.

- [48] P.B. Armstrong, R. Melchior, J.P. Quigley, Humoral immunity in long-lived arthropods, *J. Insect Physiol.* 42 (1996) 53–64.
- [49] E.A. Levashina, L.F. Moita, S. Blandin, G. Vriend, M. Lagueux, F.C. Kafatos, Conserved role of a complement-like protein in phagocytosis revealed by dsRNA knockout in cultured cells of the mosquito, *Anopheles gambiae*, *Cell* 104 (2001) 709–718.
- [50] V. Buresova, O. Hajdusek, Z. Franta, D. Sojka, P. Kopacek, IrAM-An α 2-macroglobulin from the hard tick *Ixodes ricinus*: Characterization and function in phagocytosis of a potential pathogen *Chryseobacterium indologenes*, *Dev. Comp. Immunol.* 33 (2009) 489–498.
- [51] F.F. Fagutao, M.B. Maningas, H. Kondo, T. Aoki, I. Hirano, Transglutaminase regulates immune-related genes in shrimp, *Fish. Shellfish Immunol.* 32 (2012) 711–715.
- [52] X. Lin, K. Söderhäll, I. Söderhäll, Transglutaminase activity in the hematopoietic tissue of a crustacean, *Pacifastacus leniusculus*, importance in hemocyte homeostasis, *BMC Immunol.* 9 (2008) 58.
- [53] B. Lemaitre, J. Hoffmann, The host defense of *Drosophila melanogaster*, *Annu. Rev. Immunol.* 25 (2007) 697–743.
- [54] P.H. Wang, J.P. Liang, Z.H. Gu, D.H. Wan, S.P. Weng, X.Q. Yu, et al., Molecular cloning, characterization and expression analysis of two novel Tolls (LvToll2 and LvToll3) and three putative Spatzle-like Toll ligands (LvSpz1–3) from *Litopenaeus vannamei*, *Dev. Comp. Immunol.* 36 (2012) 359–371.
- [55] F. Li, D. Wang, S. Li, H. Yan, J. Zhang, B. Wang, et al., A Dorsal homolog (FcDorsal) in the Chinese shrimp *Fenneropenaeus chinensis* is responsive to both bacteria and WSSV challenge, *Dev. Comp. Immunol.* 34 (2010) 874–883.
- [56] D. Marchant, G.K. Singhera, S. Utokaparch, T.L. Hackett, J.H. Boyd, Z. Luo, et al., Toll-like receptor 4-mediated activation of p38 mitogen-activated protein kinase is a determinant of respiratory virus entry and tropism, *J. Virol.* 84 (2010) 11359–11373.
- [57] X.Z. Shi, R.R. Zhang, Y.P. Jia, X.F. Zhao, X.Q. Yu, J.X. Wang, Identification and molecular characterization of a Spatzle-like protein from Chinese shrimp (*Fenneropenaeus chinensis*), *Fish. Shellfish Immunol.* 27 (2009) 610–617.
- [58] D.P. Parrilla-Taylor, T. Zenteno-Savín, F.J. Magallón-Barajas, Antioxidant enzyme activity in pacific whiteleg shrimp (*Litopenaeus vannamei*) in response to infection with white spot syndrome virus, *Aquaculture* 380 (2013) 41–46.
- [59] Q. Ren, R.R. Sun, X. Fan, J.X. Wang, A selenium-dependent glutathione peroxidase (Se-GPx) and two glutathione S-transferases (GSTs) from Chinese shrimp (*Fenneropenaeus chinensis*), *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 149 (2009) 613–623.
- [60] A.D. Haase, L. Jaskiewicz, H. Zhang, S. Laine, R. Sack, A. Gatignol, et al., TRBP, a regulator of cellular PKR and HIV-1 virus expression, interacts with Dicer and functions in RNA silencing, *EMBO Rep.* 6 (2005) 961–967.
- [61] S. Wang, A.J. Chen, L.J. Shi, X.F. Zhao, J.X. Wang, TRBP and eIF6 homologue in *Marsupenaeus japonicus* play crucial roles in antiviral response, *PLoS One* (2012) 7.
- [62] A. Phetrungnapha, S. Panyim, C.A. Ongvarrasopone, Tudor staphylococcal nuclease from *Penaeus monodon*: cDNA cloning and its involvement in RNA interference, *Fish. Shellfish Immunol.* 31 (2011) 373–380.