

Measuring Mass-Based Hygroscopicity of Atmospheric Particles through in Situ Imaging

Dominique S. Piens,^{*,†,@} Stephen T. Kelly,^{†,∇} Tristan H. Harder,^{†,‡} Markus D. Petters,[§] Rachel E. O'Brien,^{†,#} Bingbing Wang,^{||,⊙} Ken Teske,[⊥] Pat Dowell,[⊥] Alexander Laskin,^{||} and Mary K. Gilles^{*,†}

[†]Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States

[‡]Department of Chemistry, University of California, Berkeley, California 94720, United States


[§]Department of Marine Earth and Atmospheric Sciences, North Carolina State University, Raleigh, North Carolina 27695, United States

^{||}William R. Wiley Environmental and Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, Washington 99352, United States

[⊥]Atmospheric Radiation Monitoring (Southern Great Plains Climate Research Facility), 109596 Coal Road, Billings, Oklahoma 74630 United States

Supporting Information

ABSTRACT: Quantifying how atmospheric particles interact with water vapor is critical for understanding the effects of aerosols on climate. We present a novel method to measure the mass-based hygroscopicity of particles while characterizing their elemental and carbon functional group compositions. Since mass-based hygroscopicity is insensitive to particle geometry, it is advantageous for probing the hygroscopic behavior of atmospheric particles, which can have irregular morphologies. Combining scanning electron microscopy with energy dispersive X-ray analysis (SEM/EDX), scanning transmission X-ray microscopy (STXM) analysis, and in situ STXM humidification experiments, this method was validated using laboratory-generated, atmospherically relevant particles. Then, the hygroscopicity and elemental composition of 15 complex atmospheric particles were analyzed by leveraging quantification of C, N, and O from STXM, and complementary elemental quantification from SEM/EDX. We found three types of hygroscopic responses, and correlated high hygroscopicity with Na and Cl content. The mixing state of 158 other particles from the sample broadly agreed with those of the humidified particles, indicating the potential to infer atmospheric hygroscopic behavior from a selected subset of particles. These methods offer unique quantitative capabilities to characterize and correlate the hygroscopicity and chemistry of individual submicrometer atmospheric particles.



INTRODUCTION

Atmospheric aerosols affect the Earth's climate through direct effects, such as scattering and absorbing incident sunlight, and through indirect effects, where atmospheric particles interact with water vapor and act as cloud condensation or ice nuclei.¹ Their hygroscopic properties feedback with the uptake of volatile organics, and ultimately affect the atmospheric evolution and lifetime of particles. Quantifying how an airborne particle interacts with water vapor is critical for predicting the effect of particles on the atmospheric environment and climate, and remains an important challenge.

Multiple techniques that characterize the hygroscopic behavior of particles with precision focus on either single-particle size measurements or the effect of water vapor on an ensemble of airborne particles.^{2,3} Frequently, in field measurements, hygroscopic tandem differential mobility analysis

(HTDMA) is used to measure particle hygroscopicity.^{4–6} With HTDMA, hygroscopic growth is quantified from changes in particle size as a function of relative humidity. However, quantifying hygroscopic growth using size is susceptible to errors due to particle asphericity and particle porosity. Additionally, inferring the hygroscopic growth of internally mixed aerosols from that of their constituent compounds using size may be limited by nonideal volume additivity. Mass-based hygroscopic growth measurement provides an alternate and complementary method to quantify water uptake.

Received: February 15, 2016

Revised: April 14, 2016

Accepted: April 18, 2016

Published: April 18, 2016

The extent of water uptake, detailed microstructural changes of individual particles deposited on substrates, and correlation of microstructural changes with particle composition and/or chemistry during hydration/dehydration cycling can be investigated with spectromicroscopy. Spectromicroscopic methods include micro-FTIR,^{7,8} micro-Raman,^{9–11} micro-Raman combined with atomic force microscopy,¹² scanning and transmission electron microscopy with energy dispersive analysis of X-rays (SEM/EDX and TEM/EDX, respectively),^{13–18} and scanning transmission X-ray microscopy with near-edge X-ray absorption fine structure spectroscopy (STXM/NEXAFS).¹⁹ These are used to examine particle efflorescence and deliquescence,²⁰ changes in morphology,²¹ water vapor uptake,²² and liquid–liquid phase separation.^{23–25} STXM/NEXAFS measures optical absorption, which is proportional to mass, with high precision, both spatially and energetically resolved. STXM/NEXAFS is uniquely capable of quantitative analysis of the light elements (C, N, O) while probing their chemical bonding at spatial resolution down to 25 nm.^{19,26–28} Several groups have employed chemical imaging by STXM/NEXAFS under controlled water vapor environment.^{21,29–33} These include qualitative observations of hygroscopic properties in humic like substances (HULIS) as proxies for aerosols,³³ atmospheric particles collected from the Amazonian basin,²¹ deliquescence and efflorescence of laboratory generated particles,^{30,32} and quantitative measurements of water content in single-component proxies of atmospheric particles.^{29,34} Mikhailov et al. combined STXM/NEXAFS and electron microscopy (transmission mode) humidification experiments to investigate hygroscopicity of field-collected atmospheric aerosols.³⁵ Quantification of the mass-based hygroscopicity was done using filter samples with a differential hygroscopicity analyzer (FDHA). However, quantification of mass-based hygroscopicity for individual particles was not performed.

Here, a methodology is presented to quantify mass-based water uptake in individual submicron particles that combines STXM/NEXAFS particle characterization with in situ and water vapor uptake measurements complemented with subsequent SEM/EDX elemental composition microanalysis. The utility of in situ STXM/NEXAFS chemical imaging for mass-based water vapor uptake measurements is demonstrated using laboratory-generated particles of known compositions and uniform morphologies. Field samples, collected from the Department of Energy's Atmospheric Radiation Monitoring (DOE-ARM) site in the Southern Great Plains (SGP) field site, are then examined. First, dry characterization of the particles was done using STXM/NEXAFS, then in situ water vapor uptake experiments were performed using STXM/NEXAFS. Subsequently, SEM/EDX elemental characterization of the identical particles was performed. Finally, a composition analysis combined the STXM/NEXAFS and SEM/EDX data sets to determine mass-based hygroscopicity parameters for the atmospheric particles collected at the SGP site. This study is the first application combining STXM/NEXAFS and SEM/EDX to determine total mass of individual particles before and during hygroscopic growth both for the laboratory generated as well as field samples. The use of mass rather than size to track hygroscopic growth enabled the extension of established methods to atmospheric particles. A brief discussion on the future outlook for utilizing this analysis to extract mass-based hygroscopicity parameters from any complex heterogeneous population of atmospheric particles is presented.

■ EXPERIMENTAL SECTION

Laboratory Particle Preparation. Particles composed of NaCl, NaBr, $(\text{NH}_4)_2\text{SO}_4$, KCl, fructose, and levoglucosan were prepared by nebulizing (nebulizer, Stalder Laboratories, model 8900) a dilute solution (0.5 M) of stock chemical in deionized H_2O (18 M Ω cm resistivity). Particles then passed through a diffusion dryer, and were collected onto Si_3N_4 substrates on the seventh stage ($D_{50} = 0.56 \mu\text{m}$) of a Multi-Orifice Uniform Deposition Impactor (MOUDI; model 110-R, MSP, Inc.). To dry and recrystallize the particles, the substrates were baked at $\sim 80^\circ\text{C}$ in ambient atmosphere for several days and then stored in a desiccator until use (several days-months).

Atmospheric Particle Collection. Atmospheric particles were collected at the DOE-ARM SGP sampling site in Oklahoma, USA on March 27th 2014 at 19:00 GMT. Particle samples were collected for 20 min using a 4-stage Sioutas cascade impactor operated at a flow rate of 9 L/min. Each stage was preloaded with silicon nitride membrane substrates ($0.5 \times 0.5 \text{ mm}^2$ Si_3N_4 window size, 100 nm membrane thickness, $5 \times 5 \text{ mm}^2$ Si frame size; Silson, Inc.) and filmed TEM grids (Carbon type B film, Copper 400 mesh grids; Ted Pella, Inc.). To ensure that the samples were within the linear range of Beer's law for absorption, particles collected on the fourth stage ($D_{50} = 0.25 \mu\text{m}$) were used for microscopy analysis. After collection, the samples were placed in sealed containers, transported, and then stored in a desiccator (20–30% relative humidity) until STXM analysis (2 weeks later). Between subsequent experiments (water vapor uptake, SEM/EDX, additional STXM analysis) over the following 6 months, the samples were stored in a sealed package in a desiccator.

STXM/NEXAFS Measurements. STXM/NEXAFS was used to characterize the dry laboratory and field samples. The measurements were performed at beamline 11.0.2 of the Advanced Light Source at Lawrence Berkeley National Laboratory. Details of the STXM instrument and its operation are described elsewhere,³⁶ its applications for analysis of atmospheric particles are described in a recent review,¹⁹ and additional details on these experiments as well as nitrogen analysis are included in the [Supporting Information \(SI\) file](#). Briefly, in STXM experiments, monochromatic light from the synchrotron is focused to a small (25–40 nm) spot at the sample location by a Fresnel zone plate lens. The sample is raster scanned through the focused beam while the transmitted X-rays are collected on a single element detector behind the sample to create an image. By changing the X-ray energy, the image contrast can be changed to highlight particular molecular bonds and functional groups. Absorption at a given X-ray energy is proportional, among other factors, to the mass and composition of the matter in the X-rays' path. Collecting the same image at different X-ray energies spanning across an elemental absorption (referred to collectively as a *stack*) allows data acquisition for spatially resolved NEXAFS spectroscopy. Specific energies are then selected to acquire “short stacks” at a reduced number of energies. These “short stacks” allow more rapid mapping of carbon functional groups to allow data acquisition on a larger number of particles within the limited beamtime. Here, the number of data points collected at the C, N, and O pre-edge and postedge energies is larger than previously used.³⁷ In samples containing a significant fraction of inorganic constituents, this allows a more precise fitting of the fall off of the absorption cross section, using the Henke functions³⁸ to constrain the inorganic mass. The presence of an

element is quantitatively *mapped* by comparing an image collected either in the postedge region ($\sim 25\text{--}30$ eV beyond the elemental absorption edge for elemental mapping) or at a particular molecular resonance (to map chemical bonding) to an image collected in the pre-edge region $\sim 15\text{--}20$ eV below the energy where the element absorbs. Additional details are provided in the SI.

Characterization of Laboratory Generated Particles.

Samples of laboratory-generated particles were directly loaded in the micro reactor used for the water vapor uptake experiments. Prior to humidification, STXM/NEXAFS oxygen maps of individual dry submicron particles confirmed the absence of oxygen in NaCl, NaBr, and KCl, laboratory generated samples, indicating that they were free of water. For these salt particles as well as $(\text{NH}_4)_2\text{SO}_4$, carbon data was not acquired. For fructose and levoglucosan, both carbon short stacks and oxygen maps were collected prior to water vapor exposure.

Characterization of Particles from the Southern Great Plains. To ascertain how representative the particles used for the water vapor uptake experiments were, prior to humidification experiments, 158 additional particles from the field sample were analyzed. The STXM/NEXAFS stacks of particles from SGP field samples at the carbon K-edge provided information on their chemical composition and allowed a mixing state classification based on the amounts of organic carbon (OC), sp^2 -hybridized carbon (an indicator of soot or elemental carbon, EC), and inorganic species (IN).³⁷ For the mixing state determination, values for the optical density across the carbon edge for every pixel allowed an analysis of the chemical composition on a pixel-by-pixel basis. After analyzing the individual pixels, a compositional map of each particle was generated. Particles can contain mixtures of the three components: organic (OC), inorganic (IN), and black carbon/soot (EC). Additional analysis details are provided in Moffet et al.³⁷ The STXM particle mixing state was classified as organic carbon (OC), organic carbon with elemental carbon (OCEC), organic carbon with elemental carbon and inorganics (OCECIN) or organic carbon with inorganics (OCIN).³⁷

In Situ STXM/NEXAFS Water Vapor Uptake Measurements. The microreactor, described in Kelly et al.,³⁰ was used for in situ STXM/NEXAFS chemical imaging of particles exposed to controlled relative humidity (RH). Two helium gas flows with a combined flow rate of six standard cubic centimeters per minute, one saturated with water vapor and the other dry, were used to control RH inside the reactor. A sensor (GE Chipcap-L), located in the reactor, measured RH and temperature inside the reactor. A SiN_x window with a sample of impacted particles is mounted to the removable front metal plate using a small amount of wax (Crystalbond 509, SPI Supplies). Since the wax must be heated to be applied to the SiN_x window, impacted aerosols are exposed to a maximum temperature of up to $50\text{--}70$ °C for $15\text{--}30$ s during mounting.

After 158 particles were characterized using a higher resolution zone plate (25 nm) for inorganic content (i.e., L edges for Na, Mg, Cl, S), carbon, and oxygen, the sample was loaded in the micro reactor for the water vapor uptake experiments. Due to geometrical constraints discussed in Kelly et al.,³⁰ the water vapor uptake experiments used a 40 nm Fresnel zone plate. Visual pattern recognition was used to select three sample regions, containing particles with compositions and morphologies representative of those in the initial characterization of 158 particles. Based on previous salt

deliquescence experiments,²⁹ we selected regions with sufficient spacing between particles to limit coalescence during hygroscopic growth. For each of these regions, carbon short stacks and oxygen maps were collected under dry ($<4\%$ RH) conditions with 40 nm lateral resolution.

The RH was then increased stepwise (steps of $5\text{--}10\%$ RH) and stabilized for $5\text{--}10$ min before each set of data acquisition. At each RH step, oxygen maps (difference between absorption at the oxygen K-edge pre edge energy, 525 eV, and the post edge at 550 eV) were collected to quantify changes in the particle oxygen content attributed to water uptake. After reaching the maximum RH, the RH was decreased (steps of $10\text{--}20\%$ RH) and stabilized for $5\text{--}10$ min prior to collecting oxygen maps. The total number of single-energy images acquired on each region was approximately 120. Hence, the soft X-ray irradiation was minimized and similar to that typical of a single carbon full stack. Pattern matching with the positions and morphologies of impacted aerosols allowed rapid relocation of the identical particles for subsequent SEM/EDX analysis.

SEM/EDX Measurements. After the STXM/NEXAFS water vapor uptake experiments, the identical particles were relocated and imaged for elemental composition using SEM/EDX. An FEI Quanta digital field emission gun environmental scanning electron microscope was used in this work. The microscope is equipped with an EDAX X-ray spectrometer with a Si(Li) detector with an active area of 10 mm^2 and an ATW2 window. X-ray spectra were acquired for the same individual particles probed in the STXM experiments for 10 s, at a beam current of 430 pA and an accelerating voltage of 20 kV. Elements considered in the X-ray analysis were Na, Mg, Al, P, S, Cl, K, Ca, Mn, Fe, and Zn. Individual particle EDX spectra were preprocessed removing elements having less than 0.5 atomic %. Additional details of the SEM/EDX analysis of deposited particles are found in Laskin et al.¹⁵ and references therein.

Data Processing and Analysis. The analytical techniques, applications, and limitations of the mapping and quantitative chemical imaging of particle samples by STXM are presented elsewhere¹⁹ and only briefly discussed here. Data analysis is performed using MATLAB (Mathworks, Natick, MA). Two-dimensional transmission intensity images are obtained by raster-scanning samples of impacted particles in the STXM. In each image, the background intensity value (I_0) is determined from substrate areas without particles. I_0 is used to convert transmission intensity into optical density, (OD), using Beer's law. This analysis required additional MATLAB scripts beyond those in Moffet et al.³⁷ To identify image areas corresponding to individual particles, masks were manually drawn around particles. Individual particles at different RH values were matched visually. For each particle and RH, particle mass, m , was calculated using

$$m = \frac{\sum_i \text{OD}_i \cdot S}{\mu} \quad (1)$$

where i is a pixel in a particle's image, OD_i is the absorption at the i -th pixel at a given X-ray energy, m is the particle mass, and S is the area of a pixel. The particle's mass-absorption cross section, μ , is based on the particle's average elemental composition, and calculated using




Figure 1. Experimental (red filled triangles) and AIOMFAC theoretical (black lines) mass-based growth factors as a function of relative humidity for (a) NaCl, (b) NaBr, (c) KCl, (d) $(\text{NH}_4)_2\text{SO}_4$ labeled AS, (e) levoglucosan labeled LG, and (f) fructose. Dehydration data points are shown as green filled triangles. Deliquescence relative humidities (DRH) for NaCl and ammonium sulfate were obtained from Tang⁴⁷ and NaBr and KCl are from Cohen et al.⁴¹ The error bars show the standard error of each point with sample sizes between two and six.

$$\mu = \frac{\sum_j x_j A_j \mu_j}{\sum_j x_j A_j} \quad (2)$$

where x_j is the atomic fraction of element j , A_j is the molar mass of element j , and μ_j is the mass-absorption cross section of element j at the specific X-ray energy.³⁸ OD at the X-ray energies of an element's X-ray absorption edge and the mass-absorption cross section of that element are used in eq 1 to obtain the mass of a given element in a particle. The parameters in eq 1 and eq 2 are determined experimentally for C, N, and O using STXM/NEXAFS and using SEM/EDX for most other elements.

Dry particle masses ($\sim 4\%$ RH) and masses of the particle plus condensed water were calculated using these formulas and used to derive mass-based growth factors. This step is analogous to that employed by Ghorai and Tivanski.²⁹ For laboratory-generated particles containing oxygen, the dry mass is determined with eq 1 at the oxygen pre-edge using their known chemical formula. Laboratory-generated aerosols without oxygen (i.e., NaCl and other salts) were calculated with no dry oxygen mass (confirmed with STXM/NEXAFS imaging). To obtain the mass of the condensed water, the total oxygen mass was calculated using eq 1 at the oxygen postedge. For oxygen-containing particles (fructose, levoglucosan, and ammonium sulfate), water mass at each RH is obtained by subtracting the initial dry oxygen mass from the total oxygen mass to obtain the condensed water mass (m_w). Dry total mass (m_s) and m_w yield a mass-based growth factor at every RH (g_m).

$$g_m = \frac{m_s + m_w}{m_s} \quad (3)$$

Total Masses from STXM/NEXAFS and SEM/EDX: Atmospheric Particles of Unknown Composition. STXM/NEXAFS measurements of C, N, and O were combined with the SEM/EDX measurements for high-Z

elements to calculate the total mass of dry individual particles. In STXM/NEXAFS the sum of the OD over all pixels of a given particle is proportional to the mass-absorption cross section of the average composition of the particle, and to its total mass (eq 1). Total OD for elements other than C, N, and O is obtained by summation over a particle's STXM/NEXAFS image pixels at 278 eV (the carbon, nitrogen, and oxygen *K* pre-edges). An empirical formula for the average composition of a particle is obtained by summing its spatially resolved atomic fractions of the most common elements of Na, Mg, Al, P, S, Cl, K, Ca, Mn, Fe, and Zn measured with SEM/EDX. SEM/EDX is quantitative¹⁵ for these elements and they represent the majority of particle inorganic mass. A mass-absorption cross section, calculated from the empirical formula using eq 2, is used in eq 1 to obtain the total inorganic mass. Si was excluded because the sample substrate contains silicon. The uncertainty due to this is discussed in the SI file as well as additional information on the nitrogen content (SI Figure S1 and related discussion). Carbon, nitrogen, and oxygen masses obtained from STXM/NEXAFS were added to the inorganic mass determined with SEM/EDX to obtain the total mass of each individual dry particle.

RESULTS AND DISCUSSION

Hygroscopic Behavior of Laboratory-Generated Samples with Known Composition. In Figure 1, experimental values of the mass-based growth factor (g_m) as a function of RH obtained from STXM/NEXAFS, represented by triangles, are compared to values predicted by the thermodynamic model AIOMFAC, shown as solid black lines.^{39,40} Dashed black lines indicate the sorption isotherm of idealized crystalline salts. Prior to deliquescence, hygroscopic growth is negligible. At deliquescence, hygroscopic growth appears as a step to the g_m , as predicted by AIOMFAC. Data on the mass-based growth

factors for laboratory generated samples as a function of relative humidity are provided in the SI Tables S1–S4.

Generally, the salts (Figure 1a–d) follow idealized crystalline growth until their deliquescence RH (DRH), and after their DRH, agree with AIOMFAC. KCl (Figure 1c) and ammonium sulfate (Figure 1d) further support that trend in hygroscopic behavior because the hysteresis during their dehydration resembles the growth of aqueous particles modeled by AIOMFAC. Due to limited instrumental access, values for g_m were not obtained during dehydration for the other lab-generated samples or above DRH. The DRH observed for NaBr is consistent with those reported by Cohen et al. (between 44.5% and 45.5%)⁴¹ and Wise et al. (between 46% and 47%).¹⁸ For NaBr, the value of g_m jumps from 1.1 at 47% RH to 1.6 at 49% RH. Within the RH measurement uncertainty of 2%³⁰ the measured DRH of the crystalline salts are in agreement with literature values.

Unlike crystalline salts, which exhibit sharp deliquescence phase transition, levoglucosan and fructose exhibit gradual hygroscopic growth. Similar differences between experimental and theoretical growth factors for levoglucosan were observed by Mikhailov et al.⁴² and Mochida and Kawamura.⁴³ Measured levoglucosan g_m values increase more rapidly with higher RH than a UNIFAC-based model predicts. AIOMFAC uses UNIFAC group contribution methods to predict activity coefficients and hygroscopic behavior. Mochida and Kawamura⁴³ proposed that UNIFAC interaction coefficients were not optimized for organic molecules containing a high ratio of polar groups to nonpolar groups or closely spaced polar groups such as levoglucosan. The sorption isotherm for fructose also exhibits higher growth at high RH than predicted by AIOMFAC.

The difference between AIOMFAC and experimental values could partially be explained by a combination of our methods estimating lower initial particle mass and higher water mass at high RH than AIOMFAC. Given their continuing hygroscopic growth, sugar particles were amorphous rather than crystalline. Although the sugar particles were gently dried at 80 °C over several hours or overnight, they may have contained trace residual water and been highly viscous. Mikhailov et al. suggested that this scenario could lead to higher observed g_m values at high RH since the observed initial particle mass will be lower than the theoretical one due to water adsorption being surface limited.⁴² Therefore, differences between model assumptions and experimental conditions could explain some of the deviations in the g_m of sugars at high RH between experiment and AIOMFAC. Future experiments measuring the hygroscopic growth of lab-generated organic particles containing traces of inorganics would approximate atmospheric particles more closely, and reveal how differences between AIOMFAC and experiment impact our methods for atmospheric particles. Overall, these methods measure the hygroscopic response of individual particles of known composition in agreement with AIOMFAC.

Hygroscopic Behavior of Atmospheric Aerosol Samples of Unknown Composition. This is the first time STXM/NEXAFS and SEM/EDX are combined to determine mass-based growth factors for field collected atmospheric particles. This combination leverages the strengths of each technique and enables accurate determination of particle mass and adsorbed water mass, while also spatially identifying prominent organic chemical functional groups within individual particles.

Figure 2 depicts the range of characteristic hygroscopic curves measured for complex individual atmospheric particles




Figure 2. Mass-based growth factors of individual atmospheric particles (offset for clarity: by 1 for the green and 2 for the blue data points) grouped by their hygroscopic growth regime as a function of relative humidity. The water sorption isotherm of NaCl is overlaid on the water sorption isotherm of high hygroscopicity atmospheric aerosols.

containing both organic and inorganic components. Particles cluster into three types of hygroscopic responses labeled as low, medium, and high hygroscopicity particles. Mass-based growth factors as a function of relative humidity for these three types are provided in the SI Tables S5–S7.

The size-based growth factors of the atmospheric particles (SI Figure S2 and Table S9) yield very different growth curves. In contrast to mass-based hygroscopic growth, the particles do not cluster by type based on size growth factors obtained from their area equivalent diameter. Additionally, the 2D projections for some particles even decrease with increased water uptake, likely due to recrystallization and mechanical rearrangement as observed by Kramer et al. (1999).⁴⁴ Hence, the findings in this manuscript are only observable with its transmission imaging techniques using mass-based growth factors rather than size-based growth factors.

Complementarily, STXM/NEXAFS can quantify C, N, and O content and water uptake, while SEM/EDX can quantify inorganic elements on a single-particle basis. Figure 3 shows atomic fractions of inorganic elements for low, medium, and high hygroscopicity particles. A single correlation was found between the composition of the inorganic fraction of particles and the hygroscopic response of particles. High hygroscopicity particles have higher atomic fractions of Na and Cl than medium and low hygroscopicity particles. The range from one standard deviation below to one standard deviation above the mean Na fraction of high hygroscopicity particles is higher than and does not overlap with the same ranges for medium and low hygroscopicity particles. The trend is similar for Cl fractions, though there is some overlap between the lower range for high hygroscopicity particles and the upper range for low hygroscopicity particles. The water sorption isotherm of lab-generated NaCl is shown with the water sorption isotherm of the high hygroscopicity particles in Figure 2. The plot emphasizes that high hygroscopic growth occurs between

Particle #	Hygroscopicity	g_m (80% RH)	κ_{equiv} (80% RH)	Inorganic Atomic Fractions	4% RH	90% RH	Mixing State
1	High	2.73	0.71				OCIN
2	High	2.57	0.64				OCIN
3	High	2.01	0.41				OCECIN
4	High	1.93	0.38				OCIN
5	Medium	1.73	0.30				OCIN
6	Medium	1.45	0.18				OCECIN
7	Medium	1.34	0.14				OCIN
8	Medium	1.23	0.09				OCECIN
9	Medium	1.15	0.06				OCECIN
10	Medium	1.11	0.05				OCECIN
11	Low	1.04	0.02				OCECIN
12	Low	1.01	0.00				OCIN
13	Low	1.00	0.00				OCEC
14	Low	1.00	0.00				OCECIN
15	Low	1.00	0.00				OCECIN

0% 50% 100% 2 μm

Figure 3. Table presenting hygroscopicity, mass-based growth factor at 80% RH (g_m), hygroscopicity parameter κ_{equiv} at 80% RH, inorganic atomic composition, STXM images at 4% and 90% RH acquired at 525 eV, and mixing state classification for 15 atmospheric particles.




Figure 4. Left: STXM based mixing state distribution of all particles characterized on the Southern Great Plains sample (158) and the subset of the 15 particles used during water vapor uptake experiments. Mixtures of individual components (organic (OC), inorganic (IN), and black carbon/soot (EC)) within individual particles are indicated. Error bars calculated by varying the thresholds for classifying individual components by 10% are shown. Right: the corresponding size distributions. The error bars represent the change in size range frequencies due to an estimated uncertainty of ± 2 pixels on area equivalent diameter computation.

72% and 78% RH for high hygroscopicity particles, very near NaCl DRH (75.3% RH). Laboratory-generated NaCl and high hygroscopicity particles have similar discontinuities in hygro-

scopic growth. This suggests that Na and Cl in the form of NaCl may drive the hygroscopic growth of high hygroscopicity particles. Atomic fractions for the other elements show

significant overlap between particles with different hygroscopicities.

To provide an alternate representation of the hygroscopicity of the atmospheric particles, another parameter representing aerosol-water interactions is presented in Figure 3. Petters and Kreidenweis introduced the use of a single parameter, κ , to represent the effect a particle has on water activity by calculating the ratio of the volume of water to particulate matter.⁴² Higher values of κ correspond to higher hygroscopicity and for atmospherically relevant particles κ typically ranges from 0 to values exceeding 1. However, instead of a ratio of volumes mass-based growth factors can also be used to calculate κ by assuming particle density. A density-equivalent κ_{equiv} , which is defined as the κ value of a particle with the observed hygroscopic mass growth factor and assumed dry density, ρ_{d} , is calculated with eq 4:

$$\kappa_{\text{equiv}} = (a_{\text{w}}^{-1} - 1)(g_{\text{m}} - 1)r \quad (4)$$

where r is the ratio of dry particle density to the density of pure water. Here we assume $r = 1.760$ based on the density of solid ammonium sulfate. Kelvin effects are neglected and water activity is taken to be the equilibrium RH. Density equivalent κ provides an approximate comparison to volume-growth factor published data in the literature.^{42–44} If the studied particle density equals that of assumed value, the reported κ_{equiv} is identical to κ obtained via volumetric methods. A 10% relative difference between actual and assumed particle density results in a 10% difference to values inferred from the volumetric-based method. The density equivalent κ_{equiv} values can be compared to the known κ values of other substances.^{42–44} In this study, the most hygroscopic atmospheric particle observed has a density equivalent κ_{equiv} of 0.71 at high RH (80%), which is much lower than the estimated κ_{equiv} value reported for NaCl, but near the estimated κ_{equiv} values for most sulfate and nitrate salts typically found in atmospheric aerosols.⁴⁵ Medium and low hygroscopicity particles have κ_{equiv} values in the same range as organic particles, attaining the lowest possible value of 0 for the low hygroscopicity particles.⁴⁶

As described above, STXM/NEXAFS at the carbon K-edge identifies three components: organic (OC), inorganic (IN), and black carbon/soot (EC) to define a mixing state.³⁷ SI Figure S3 contains images that illustrate how the mixing state is determined for particles and SI Table S8 provides details on the mixing state classification. Figure 4 (left panel) shows the STXM based mixing states of the particles used for hygroscopic growth experiments (green) compared to that of the sample of 158 particles (blue). Based upon a visual analysis of preliminary STXM images we anticipated that these 15 particles were reasonably representative of the samples. Hence, they were selected for water vapor uptake experiments. Figure 4 (right panel) compares the size distribution of the 158 particles characterized using STXM/NEXAFS (blue) compared to that of the 15 particles selected for water vapor uptake experiments (green). Overall, the particle size distributions match well between the two sample sets. The agreement for the mixing state is lower (Figure 4, left panel) indicating that field samples containing more diverse particle-types would require better statistical depth. With the sample size of 15 particles, no significant correlation was found between hygroscopicity, and mixing state or particle size. However, if correlations were observed, by using a statistically significant subset of the population, then estimates for the hygroscopic behavior of the

sample could be obtained. Additional characterization of the IN, via analysis with SEM/EDX, would be required as kappa values vary for different inorganic compounds.⁴⁶

STXM/NEXAFS has previously been used to quantify the hygroscopicity of laboratory generated particles of known composition^{29,30,32,33} and to observe the hygroscopic behavior of ambient particles of unknown composition.²¹ Here, we demonstrated that the combination of STXM/NEXAFS and SEM/EDX could quantitatively determine the hygroscopic behavior of both homogeneous particles with known composition and mixed atmospheric particles of unknown composition. Mass-based growth factors of mixed atmospheric particles of unknown composition were obtained on a per-particle basis by combining SEM/EDX measurements with STXM/NEXAFS characterizations. This method was validated using laboratory surrogates. Such data on field samples may provide means to correlate hygroscopic behavior and particle composition and chemistry. Although we attempted to find correlations between carbon functional groups (C=C, R-OH, COOH) and particle hygroscopicity, no apparent relationship was found. The mass-based growth factors can be used to characterize the statistics and range of hygroscopic behavior present in a sample of collected atmospheric particles and relate this range of behavior to elemental and functional chemistry.

■ ASSOCIATED CONTENT

📄 Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.est.6b00793.

More detailed experimental methods for STXM/NEXAFS measurements and the characterization of SGP samples; the effects of uncertainty in nitrogen and silicon measurements on the mass-based growth factors of the SGP atmospheric particles; plot of size-based growth factors as a function of RH for the SGP atmospheric particles; tables containing the experimental data displayed in Figures 1–4; images illustrating the mixing state of particles in terms of OC, EC and IN (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*(D.S.P.) E-mail: dpiens@stanford.edu.

*(M.K.G.) E-mail: mkgilles@lbl.gov; phone: 510-495-2775.

Present Addresses

@(D.S.P.) Stanford University, Stanford, California 94305, United States.

∇(S.T.K.) Carl Zeiss X-ray Microscopy Inc., Pleasanton, California 94588, United States.

#(R.E.O.) Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States.

⊙(B.W.) State Key Laboratory of Marine Environmental Science, College of Ocean and Earth Sciences, Xiamen University, Xiamen 361102, China.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

STXM/NEXAFS dry characterization experiments (MKG, STK, THH) were supported by the Atmospheric Systems Research (ASR) of the U.S. Department of Energy (DOE),

Office of Science, Office of Biological and Environmental Research (OBER) program. Dry STXM/NEXAFS analysis of the field samples (T.H.H.) was partially supported by the German National Academic Foundation. Development of the MATLAB scripts and analysis for the mass-based water vapor uptake analysis (D.S.P.) was supported through the DOE, Office of Workforce Development for Teachers and Scientists (WDTs) under the Science Undergraduate Laboratory Internship (SULI) program. R.E.O. (supported by DOE ASR) provided analysis guidance to both D.S.P. and T.H.H. The experimental portion of the water vapor uptake experiments (S.T.K., M.K.G.) and assistance with the interpretation of the data (M.P.) was supported by the ASR DOE OBER program. STXM/NEXAFS measurements were done with the assistance of Tolek Tyliczszak at Beamline 11.0.2 of the Advanced Light Source (ALS) at Lawrence Berkeley National Laboratory (LBNL). The ALS is supported by the Director, Office of Science, Office of Basic Energy Sciences (BES), of the U.S. DOE under Contract No. DE-AC02-05CH11231. Beamline 11.0.2 is supported under the same contract as well as the Condensed Phase and Interfacial Molecular Sciences Program of DOE BES. A.L. and B.W. acknowledge support from the Chemical Imaging Initiative of the Laboratory Directed Research and Development program at Pacific Northwest National Laboratory (PNNL). The preparation of laboratory samples and SEM/EDX measurements were performed at the W. R. Wiley Environmental Molecular Sciences Laboratory (EMSL) - a national scientific user facility located at PNNL, and sponsored by DOE OBER. PNNL is operated for U.S. DOE by Battelle Memorial Institute under Contract No. DE-AC06-76RL0 1830. Samples were collected (K.T., P.D.) at the Atmospheric Radiation Monitoring (ARM) site located at Southern Great Plains and sponsored by DOE OBER.

REFERENCES

- (1) Andreae, M. O.; Rosenfeld, D. Aerosol-cloud-precipitation interactions. Part I. The nature and sources of cloud-active aerosols. *Earth-Sci. Rev.* **2008**, *89* (1–2), 13–41.
- (2) Nguyen, T. K. V.; Petters, M. D.; Suda, S. R.; Guo, H.; Weber, R. J.; Carlton, A. G. Trends in particle-phase liquid water during the Southern Oxidant and Aerosol Study. *Atmos. Chem. Phys.* **2014**, *14* (20), 10911–10930.
- (3) Swietlicki, E.; Hansson, H. C.; Hameri, K.; Svenningsson, B.; Massling, A.; McFiggans, G.; McMurry, P. H.; Petaja, T.; Tunved, P.; Gysel, M.; Topping, D.; Weingartner, E.; Baltensperger, U.; Rissler, J.; Wiedensohler, A.; Kulmala, M. Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments - a review. *Tellus, Ser. B* **2008**, *60* (3), 432–469.
- (4) Suda, S. R.; Petters, M. D. Accurate determination of aerosol activity coefficients at relative humidities up to 99% using the hygroscopicity tandem differential mobility analyzer technique. *Aerosol Sci. Technol.* **2013**, *47* (9), 991–1000.
- (5) Yeung, M. C.; Lee, B. P.; Li, Y. J.; Chan, C. K. Simultaneous HTDMA and HR-ToF-AMS measurements at the HKUST Supersite in Hong Kong in 2011. *J. Geophys. Res.-Atmos.* **2014**, *119* (16), 9864–9883.
- (6) Martin, S. T.; Rosenoern, T.; Chen, Q.; Collins, D. R. Phase changes of ambient particles in the Southern Great Plains of Oklahoma. *Geophys. Res. Lett.* **2008**, *35* (22), L22801.
- (7) Liu, Y.; Laskin, A. Hygroscopic properties of $\text{CH}_3\text{SO}_3\text{Na}$, $\text{CH}_3\text{SO}_3\text{NH}_4$, $(\text{CH}_3\text{SO}_3)_2\text{Mg}$, and $(\text{CH}_3\text{SO}_3)_2\text{Ca}$ particles studied by micro-FTIR spectroscopy. *J. Phys. Chem. A* **2009**, *113* (8), 1531–1538.
- (8) Liu, Y.; Yang, Z.; Desyaterik, Y.; Gassman, P. L.; Wang, H.; Laskin, A. Hygroscopic behavior of substrate-deposited particles studied by micro-FT-IR spectroscopy and complementary methods of particle analysis. *Anal. Chem.* **2008**, *80* (3), 633–642.
- (9) Ciobanu, V. G.; Marcolli, C.; Krieger, U. K.; Weers, U.; Peter, T. Liquid-liquid phase separation in mixed organic/inorganic aerosol particles. *J. Phys. Chem. A* **2009**, *113* (41), 10966–10978.
- (10) Yeung, M. C.; Chan, C. K. Water content and phase transitions in particles of inorganic and organic species and their mixtures using micro-Raman spectroscopy. *Aerosol Sci. Technol.* **2010**, *44* (4), 269–280.
- (11) Garland, R. M.; Wise, M. E.; Beaver, M. R.; DeWitt, H. L.; Aiken, A. C.; Jimenez, J. L.; Tolbert, M. A. Impact of palmitic acid coating on the water uptake and loss of ammonium sulfate particles. *Atmos. Chem. Phys.* **2005**, *5*, 1951–1961.
- (12) Laskina, O.; Morris, H. S.; Grandquist, J. R.; Qin, Z.; Stone, E. A.; Tivanski, A. V.; Grassian, V. H. Size matters in the water uptake and hygroscopic growth of atmospherically relevant multicomponent aerosol particles. *J. Phys. Chem. A* **2015**, *119* (19), 4489–97.
- (13) Laskin, A. Electron beam analysis and microscopy of individual particles. In *Fundamentals and Applications in Aerosol Spectroscopy*; Signorell, R., Reid, J. P., Eds.; CRC Press Taylor & Francis: Boca Raton, FL, 2010; pp 463–491.
- (14) Posfai, M.; Buseck, P. R. Nature and climate effects of individual tropospheric aerosol particles. *Annu. Rev. Earth Planet. Sci.* **2010**, *38*, 17–43.
- (15) Laskin, A.; Cowin, J. P.; Iedema, M. J. Analysis of individual environmental particles using modern methods of electron microscopy and X-ray microanalysis. *J. Electron Spectrosc. Relat. Phenom.* **2006**, *150* (2–3), 260–274.
- (16) Ahn, K. H.; Kim, S. M.; Jung, H. J.; Lee, M. J.; Eom, H. J.; Maskey, S.; Ro, C. U. Combined use of optical and electron microscopic techniques for the measurement of hygroscopic property, chemical composition, and morphology of individual aerosol particles. *Anal. Chem.* **2010**, *82* (19), 7999–8009.
- (17) Eom, H. J.; Gupta, D.; Li, X.; Jung, H. J.; Kim, H.; Ro, C. U. Influence of collecting substrates on the characterization of hygroscopic properties of inorganic aerosol particles. *Anal. Chem.* **2014**, *86* (5), 2648–56.
- (18) Wise, M. E.; Biskos, G.; Martin, S. T.; Russell, L. M.; Buseck, P. R. Phase transitions of single salt particles studied using a transmission electron microscope with an environmental cell. *Aerosol Sci. Technol.* **2005**, *39* (9), 849–856.
- (19) Moffet, R. C.; Tivanski, A. V.; Gilles, M. K., Scanning transmission X-ray microscopy: Applications in atmospheric aerosol research. In *Fundamentals and Applications in Aerosol Spectroscopy*; Signorell, R., Reid, J. P., Eds.; CRC Press Taylor & Francis: Boca Raton, FL, 2010; pp 420–462.
- (20) Freney, E. J.; Martin, S. T.; Buseck, P. R. Deliquescence and efflorescence of potassium salts relevant to biomass-burning aerosol particles. *Aerosol Sci. Technol.* **2009**, *43* (8), 799–807.
- (21) Pöhlker, C.; Saturno, J.; Kruger, M. L.; Forster, J. D.; Weigand, M.; Wiedemann, K. T.; Bechtel, M.; Artaxo, P.; Andreae, M. O. Efflorescence upon humidification? X-ray microspectroscopic in situ observation of changes in aerosol microstructure and phase state upon hydration. *Geophys. Res. Lett.* **2014**, *41* (10), 3681–3689.
- (22) Ghorai, S.; Wang, B. B.; Tivanski, A.; Laskin, A. Hygroscopic properties of internally mixed particles composed of NaCl and water-soluble organic acids. *Environ. Sci. Technol.* **2014**, *48* (4), 2234–2241.
- (23) You, Y.; Renbaum-Wolff, L.; Carreras-Sospedra, M.; Hanna, S. J.; Hiranuma, N.; Kamal, S.; Smith, M. L.; Zhang, X. L.; Weber, R. J.; Shilling, J. E.; Dabdub, D.; Martin, S. T.; Bertram, A. K. Images reveal that atmospheric particles can undergo liquid-liquid phase separations. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109* (33), 13188–13193.
- (24) You, Y.; Smith, M. L.; Song, M. J.; Martin, S. T.; Bertram, A. K. Liquid-liquid phase separation in atmospherically relevant particles consisting of organic species and inorganic salts. *Int. Rev. Phys. Chem.* **2014**, *33* (1), 43–77.
- (25) O'Brien, R. E.; Wang, B.; Kelly, S. T.; Lundt, N.; You, Y.; Bertram, A. K.; Laskin, A.; Gilles, M. K. Imaging of liquid-liquid phase

separation in aerosol particles at nanometer scale. *Environ. Sci. Technol.* **2015**, *49*, 4995–5002.

(26) Tivanski, A. V.; Hopkins, R. J.; Tyliczszak, T.; Gilles, M. K. Oxygenated interface on biomass burn tar balls determined by single particle scanning transmission X-ray microscopy. *J. Phys. Chem. A* **2007**, *111* (25), 5448–5458.

(27) Hopkins, R. J.; Tivanski, A. V.; Marten, B. D.; Gilles, M. K. Chemical bonding and structure of black carbon reference materials and individual carbonaceous atmospheric aerosols. *J. Aerosol Sci.* **2007**, *38* (6), 573–591.

(28) Maria, S. F.; Russell, L. M.; Gilles, M. K.; Myneni, S. C. B. Organic aerosol growth mechanisms and their climate-forcing implications. *Science* **2004**, *306* (5703), 1921–1924.

(29) Ghorai, S.; Tivanski, A. V. Hygroscopic behavior of individual submicrometer particles studied by X-ray spectromicroscopy. *Anal. Chem.* **2010**, *82* (22), 9289–9298.

(30) Kelly, S. T.; Nigge, P.; Prakash, S.; Laskin, A.; Wang, B.; Tyliczszak, T.; Leone, S. R.; Gilles, M. K. An environmental sample chamber for reliable scanning transmission X-ray microscopy measurements under water vapor. *Rev. Sci. Instrum.* **2013**, *84*, 073708.

(31) Steimer, S. S.; Lampimaki, M.; Coz, E.; Grzanic, G.; Ammann, M. The influence of physical state on shikimic acid ozonolysis: A case for in situ microspectroscopy. *Atmos. Chem. Phys.* **2014**, *14* (19), 10761–10772.

(32) Zelenay, V.; Ammann, M.; Krepelova, A.; Birrer, M.; Tzvetkov, G.; Vernooij, M. G. C.; Raabe, J.; Huthwelker, T. Direct observation of water uptake and release in individual submicrometer sized ammonium sulfate and ammonium sulfate/adipic acid particles using X-ray microspectroscopy. *J. Aerosol Sci.* **2011**, *42* (1), 38–51.

(33) Zelenay, V.; Huthwelker, T.; Krepelova, A.; Rudich, Y.; Ammann, M. Humidity driven nanoscale chemical separation in complex organic matter. *Environ. Chem.* **2011**, *8* (4), 450–460.

(34) Ghorai, S.; Laskin, A.; Tivanski, A. V. Spectroscopic evidence of keto-enol tautomerism in deliquesced malonic acid particles. *J. Phys. Chem. A* **2011**, *115* (17), 4373–4380.

(35) Mikhailov, E. F.; Mironov, G. N.; Pohlker, C.; Chi, X.; Kruger, M. L.; Shiraiwa, M.; Forster, J. D.; Poschl, U.; Vlasenko, S. S.; Ryskhovich, T. I.; Weigand, M.; Kilcoyne, A. L. D.; Andreae, M. O. Chemical composition, microstructure, and hygroscopic properties of aerosol particles at the Zotino Tall Tower Observatory (ZOTTO), Siberia, during a summer campaign. *Atmos. Chem. Phys.* **2015**, *15* (15), 8847–8869.

(36) Kilcoyne, A. L. D.; Tyliczszak, T.; Steele, W. F.; Fakra, S.; Hitchcock, P.; Franck, K.; Anderson, E.; Harteneck, B.; Rightor, E. G.; Mitchell, G. E.; Hitchcock, A. P.; Yang, L.; Warwick, T.; Ade, H. Interferometer-controlled scanning transmission X-ray microscopes at the Advanced Light Source. *J. Synchrotron Radiat.* **2003**, *10*, 125–136.

(37) Moffet, R. C.; Henn, T.; Laskin, A.; Gilles, M. K. Automated chemical analysis of internally mixed aerosol particles using X-ray spectromicroscopy at the carbon K-edge. *Anal. Chem.* **2010**, *82* (19), 7906–7914.

(38) Henke, B. L.; Gullikson, E. M.; Davis, J. C. X-ray interactions -photoabsorption, scattering, transmission, and reflection at $E = 50$ – $30,000$ eV, $Z = 1$ – 92 . *At. Data Nucl. Data Tables* **1993**, *54* (2), 181–342.

(39) Zuend, A.; Marcolli, C.; Booth, A. M.; Lienhard, D. M.; Soonsin, V.; Krieger, U. K.; Topping, D. O.; McFiggans, G.; Peter, T.; Seinfeld, J. H. New and extended parameterization of the thermodynamic model AIOMFAC: calculation of activity coefficients for organic-inorganic mixtures containing carboxyl, hydroxyl, carbonyl, ether, ester, alkenyl, alkyl, and aromatic functional groups. *Atmos. Chem. Phys.* **2011**, *11* (17), 9155–9206.

(40) Zuend, A.; Marcolli, C.; Luo, B. P.; Peter, T. A thermodynamic model of mixed organic-inorganic aerosols to predict activity coefficients. *Atmos. Chem. Phys.* **2008**, *8* (16), 4559–4593.

(41) Cohen, M. D.; Flagan, R. C.; Seinfeld, J. H. Studies of concentrated electrolyte-solutions using the electrodynamic balance 0.3. Solute nucleation. *J. Phys. Chem.* **1987**, *91* (17), 4583–4590.

(42) Mikhailov, E.; Vlasenko, S.; Martin, S. T.; Koop, T.; Pöschl, U. Amorphous and crystalline aerosol particles interacting with water vapor: conceptual framework and experimental evidence for restructuring, phase transitions and kinetic limitations. *Atmos. Chem. Phys.* **2009**, *9* (24), 9491–9522.

(43) Mochida, M.; Kawamura, K. Hygroscopic properties of levoglucosan and related organic compounds characteristic to biomass burning aerosol particles. *J. Geophys. Res.-Atmos.* **2004**, *109* (D21), D21202.

(44) Kramer, L.; Poschl, U.; Niessner, R. Microstructural rearrangement of sodium chloride condensation aerosol particles on interaction with water vapor. *J. Aerosol Sci.* **2000**, *31* (6), 673–685.

(45) Kreidenweis, S. M.; Petters, M. D.; DeMott, P. J. Single-parameter estimates of aerosol water content. *Environ. Res. Lett.* **2008**, *3* (3), 035002.

(46) Petters, M. D.; Kreidenweis, S. M. A single parameter representation of hygroscopic growth and cloud condensation nucleus activity. *Atmos. Chem. Phys.* **2007**, *7* (8), 1961–1971.

(47) Tang, I. N. Thermodynamic and optical properties of mixed-salt aerosols of atmospheric importance. *J. Geophys. Res.-Atmos.* **1997**, *102* (D2), 1883–1893.