



# Fishing for feed in China: Facts, impacts and implications

Wenbo Zhang<sup>1</sup> | Min Liu<sup>2</sup> | Yvonne Sadovy de Mitcheson<sup>3</sup> | Ling Cao<sup>4,5</sup> |  
Duncan Leadbitter<sup>6</sup> | Richard Newton<sup>7</sup> | David C. Little<sup>7</sup> | Songlin Li<sup>1</sup> | Yi Yang<sup>8</sup> |  
Xiao Chen<sup>9</sup> | Wei Zhou<sup>8</sup>

<sup>1</sup>National Demonstration Center for Experimental Fisheries Science Education, Shanghai Engineering Research Center of Aquaculture, Centre for Research on Environmental Ecology and Fish Nutrition of the Ministry of Agriculture, Shanghai Ocean University, Shanghai, China

<sup>2</sup>State Key Laboratory of Marine Environmental Science and College of Ocean & Earth Sciences, Xiamen University, Xiamen, Fujian, China

<sup>3</sup>School of Biological Sciences, Swire Institute of Marine Science, University of Hong Kong, Hong Kong, China

<sup>4</sup>Institute of Oceanography, Shanghai Jiao Tong University, Shanghai, China

<sup>5</sup>Center on Food Security and the Environment, Stanford University, Stanford, CA, USA

<sup>6</sup>Australian National Centre for Ocean Resources and Security, University of Wollongong, Wollongong, NSW, Australia

<sup>7</sup>Institute of Aquaculture, University of Stirling, Stirling, UK

<sup>8</sup>Greenpeace East Asia, Beijing, China

<sup>9</sup>College of Marine Science, South China Agricultural University, Guangzhou, China

## Correspondence

Wenbo Zhang, National Demonstration Center for Experimental Fisheries Science Education, Shanghai Ocean University, Shanghai 201306, China.  
Email: wb-zhang@shou.edu.cn

Min Liu, State Key Laboratory of Marine Environmental Science and College of Ocean & Earth Sciences, Xiamen University, Xiamen 361102, Fujian, China.  
Email: minliuxm@xmu.edu.cn

Wei Zhou, Greenpeace East Asia, Beijing 100007, China.  
Email: zhou.wei@greenpeace.org

## Funding information

China-ASEAN Maritime Cooperation Fund; National Natural Science Foundation of China, Grant/Award Number: 41476149; State Oceanic Administration of China, Grant/Award Number: 220203993022761133; ADM Capital Foundation; Swire Trust

## Abstract

China is the world's largest capture fisheries and aquaculture producer. Over recent decades, China's domestic marine catch composition has changed markedly, from large volumes of a few high-valued food species to multiple, small, low-valued, species, a significant proportion of which is primarily used as animal, especially fish, feed. Despite the growing volume and economic importance of the feed catches, their species composition, catch volumes and socio-environmental impacts are all poorly understood. Based on a nationwide survey of >800 fishing vessels, and the identification and measurement of >12,000 fish and invertebrate individuals, the present study provides an overview of the feed component of China's domestic marine catch, by volumes, species and sizes, and found it to be substantial and biologically unsustainable. Half of the trawler catch (3 million metric tons, mmt), or 35% of the total catch (4.6 mmt) in China's exclusive economic zone, are now comprised of low-valued "feed-grade fish". The present study identified 218 fish species, 50 crustaceans and five cephalopods, and of these, 102 fish species were food species with 89% individuals in their juvenile size ranges. Feed-grade fish were mainly used as aquaculture feed directly, or indirectly, through the feed industry after reduction to fishmeal and fish oil. The unparalleled scale and poor fisheries resource condition of China's domestic marine fisheries, in parallel with severe overfishing of juveniles, creates a demand for fundamental changes to fishery management practices, including a significant reduction of fishing effort to ensure productivity and ecosystem resilience.

Wenbo Zhang and Min Liu contributed equally.

## KEYWORDS

biodiversity, feed-grade fish, management, multispecies fisheries, trash fish, trawl

## 1 | INTRODUCTION

China (mainland, excluding Hong Kong, Macau and Taiwan unless otherwise specified) is the world's largest capture fisheries and aquaculture producing country, accounting for 19.2% of global marine capture fisheries production and 61.5% of global aquaculture production in 2016 (FAO, 2018). In 2016, China's total domestic marine capture fisheries production was 13.3 mmt (million metric tons), and total aquaculture (marine and freshwater) production was 51.4 mmt (MOA, 2017a). Capture fisheries and aquaculture in China provide food and income, 13.8 million jobs, and livelihoods for nearly 20 million people (MOA, 2017a).

Despite the introduction of a range of marine fisheries management measures at both national and local levels since the 1980s (Cao et al., 2017), fishing pressure has grown over the last four decades (Figure S1). Persistently high fishing pressure, together with other pervasive deleterious anthropogenic influences, such as runoff of fertilizers, habitat loss and degradation, has resulted in sharp declines in fishery biomass and catch per unit effort (CPUE) in China's major fishing grounds. In the Bohai Sea, the CPUE declined from >420 kg haul<sup>-1</sup> hr<sup>-1</sup> in 1959, to ~160 kg haul<sup>-1</sup> hr<sup>-1</sup> in 1982, ~30 kg haul<sup>-1</sup> hr<sup>-1</sup> in 1993, and <8 kg haul<sup>-1</sup> hr<sup>-1</sup> in 1998–2011 (Jin et al., 2013; Shan, Sun, Jin, Li, & Dai, 2013). In the Yellow Sea and East China Sea, the CPUE dropped from 73.54 to 39.19 kg haul<sup>-1</sup> hr<sup>-1</sup> in just ten years between 1991 and 2000 (Cheng & Yu, 2004). In the northern South China Sea, the fishery biomass declined from 2.7 mt km<sup>-2</sup> in the 1930s, to 1.5 mt km<sup>-2</sup> in 1956, 1.1 mt km<sup>-2</sup> in 1960–1973, and 0.3 mt km<sup>-2</sup> in 1997–1999 (Qiu, 2002).

In concert with biomass and CPUE declines, the species composition of China's domestic marine catches over the past six decades has undergone substantial changes from a predominance of a few medium to relatively large-sized, high-valued, high trophic level, demersal species to multiple small, low-valued, short-lived, low trophic level (Cao et al., 2017; Liu & Sadovy de Mitcheson, 2008; Shen & Heino, 2014) (Figure S2). These changes are largely attributable to the overfishing of these larger species driven by ineffective management resulting in increased numbers of smaller, lower trophic level species (Cao et al., 2017; Lundgren, Staples, Funge-Smith, & Clausen, 2006; Pauly, Christensen, Dalsgaard, Froese, & Torres, 1998; Shen & Heino, 2014).

Since the 1990s, fisheries production has increasingly been used to supply fishmeal and animal feed, particularly aquaculture feed, shifting away from its historic use for direct human consumption. This pattern is observed not only in China (Cao et al., 2015; Funge-Smith, Lindebo, & Staples, 2005; Lundgren et al., 2006), but also in Vietnam (Edwards, Tuan, & Allan, 2004) and several other Asian countries and areas (Funge-Smith et al., 2005). The increasing importance of animal feed during a period when food fisheries were

1. INTRODUCTION	48
2. METHODOLOGY	49
2.1. Terms used	50
2.2. Fishing vessel/gear survey	50
2.3. Fish sampling, identification and measurement	50
2.4. Data analysis	51
3. RESULTS	51
3.1. Proportion of feed-grade fish in catch	51
3.2. Feed-grade fish production	51
3.3. Use and prices of feed-grade fish	52
3.4. Species composition	52
3.5. Food fish species and proportion of catches composed of juveniles	53
3.6. Stock assessment and conservation status	53
4. DISCUSSION	54
4.1. Fishing for feed in China is substantial and biologically unsustainable	54
4.2. Effective management is needed for both food and feed components	55
5. CONCLUSIONS	59
ACKNOWLEDGEMENTS	60
CONFLICT OF INTEREST	60
DATA AVAILABILITY STATEMENT	60
REFERENCES	60

declining enabled some fisheries to remain viable; for example in Hong Kong after declines in traditional trawl fisheries occurred due to overfishing, fishers were increasingly able to earn money from smaller, poorer quality fish, being sold as animal feed for the burgeoning mariculture sector. (Cheung & Sadovy, 2004).

Due to the highly diverse nature of the world's fisheries and because of different terminologies and histories among countries (FAO, 2011b), there is no single or consistent definition or term for fisheries fishing for feed, or their product, that clearly describes the specific proportion of the catch that is used as aquaculture (or live-stock) feed. In the past, when fisheries were in good condition and aquaculture less developed, small-sized, low-valued fish were typically discarded and variously referred to as "discards" or "bycatch" or "trash fish," reflecting the fact that they were not targeted, were incidental or were of little or no economic value, and often being unfit for human consumption due to small size and/or poor condition (Alverson, Freeberg, Murawski, & Pope, 1994; FAO, 2011a; Nguyen et al., 2009). In the early days of fishery development, this discarding

in Southeast Asia was responsible for an estimated loss of some 3.4 mmt of fish catch per year (Chee, 1997).

Other terms have been variously and inconsistently used, making comparisons across studies a challenge. Terms for the catch component not directly used for human food initially reflected what happened to the catch while more recently terms refer to, the fisheries that target fish used as feed. Catch terms range from “pig fish” (Edwards et al., 2004; Funge-Smith et al., 2005), “low value/trash fish” (Funge-Smith et al., 2005; Lundgren et al., 2006), “mixed fish feed” (Chau & Sadovy, 2005) to “feed fish” (Huntington & Hasan, 2009). Although reduction fisheries have existed for centuries for industrial purposes (Smith, 1991), as the use of catches for animal feed became more important, specific fisheries terminology began to be used, such as “feed fisheries” (Hasan & Halwart, 2009) and “forage fisheries” (Watson, Alder, & Pauly, 2006), which were used alongside the term “reduction fisheries” (Parker & Tyedmers, 2011; Shepherd & Jackson, 2013).

Some terms for this “feed” component fisheries are no longer fit for purpose or are even misleading. The best example is the still widely used term “trash fish”. As fisheries focused on food fish declined and aquaculture increased, a growing proportion of “trash fish” was retained and came to have economic value in the aquaculture (or other livestock) sector (Alverson et al., 1994; Funge-Smith et al., 2005; Zeller, Cashion, Palomares, & Pauly, 2018). Moreover, what may be by-catch in one fishery or in one season, may be valuable food fish in another or would become food fish if animals were retained in better condition or were taken at larger body sizes. Along with the development of aquatic food processing technologies and improvements in the handling of fish, there was also a rapid expansion of markets for other human food products such as surimi, fish balls, sauces or pastes. Species once used as by-catch but suited for processing into surimi are increasingly targeted or diverted from the fishmeal supply chain and used as raw material (Anon, 2016; Funge-Smith, Briggs, & Miao, 2012; Siriraksophon, Pangsorn, & Laong-manee, 2009).

While in many parts of the world the “feed” component is primarily (if not solely) caught from targeted forage (or reduction) fisheries (FAO, 2011a; Hasan & Halwart, 2009), the sourcing of it in China is more complicated. This is because multiple fishing gears are involved which take a complex mix of species that are both targeted and non-targeted and can be used as direct feed for aquatic animals and, indirectly, to produce fishmeal, or for direct human consumption, according to their condition, species and size. In the present study, we use the term “feed-grade fish” to describe those fish (and invertebrates) that are targeted or retained to be used as fish or livestock feed.

Over the past two decades, fishing for feed-grade fish in China has become increasingly important, due to overfishing of food fish associated with poor fisheries management, and, growth in demand for feed-grade fish, especially with the development of aquaculture (Cao et al., 2015; Funge-Smith et al., 2005; Huntington & Hasan, 2009; Naylor et al., 2000). Into the new millennium, China's aquaculture has been greatly intensified with much more formulated feed and direct feeding of feed-grade fish required (Chiu et al., 2013; Edwards, 2008a, 2008b). There has been a trend to “farming up” the

food web with increasingly more high value species, especially carnivores, being farmed (Miao & Liao, 2007; Naylor et al., 2000; Zhou, 2007), further contributing to the increasing demand for feed-grade fish.

The feed component is today a significant aspect of China's domestic marine fisheries. Estimates based on FAO data over a decade ago show that the feed component represented more than one-third of China's marine catch (FAO, 2007; Funge-Smith et al., 2005; Han & Xu, 2004) and was mainly constituted by unidentified “nei” (not elsewhere included) species and forage species (Cao et al., 2015; Han & Xu, 2004). Sporadic surveys also suggest that 26%–90% of the fisheries catch by weight or number in China's EEZ are feed-grade fish, especially from non-targeted trawls and stow nets, in some cases higher than the situation found in other Southeast Asian countries (Table S1) but similar to countries where targeted reduction fisheries are operating.

However, the feed component of China's domestic marine fisheries has not been documented systemically at the national level. Its scale, species composition, as well as biological, ecosystem and management implications are poorly understood. For example, “nei” accounts for approximately 31% of China's domestic marine catch (Cao et al., 2015), but its composition is largely unknown. This situation makes it difficult to assess the impact of the source fisheries and seriously constrains the development of effective management.

Meanwhile, there are growing concerns that the demonstrable declines in large-scale, and unmanaged, fisheries bring substantial risks to the marine ecosystem, threaten biodiversity and endanger long-term marine ecosystem resilience. One critical risk is that further overfishing of a wide range of species and sizes in China's EEZ will seriously reduce the future productivity of marine fisheries (Cao et al., 2017, 2015).

The objectives of the present study were to reveal the complicated nature of the feed component of China's domestic marine catch, related fisheries, volumes, species composition, destination/use of catches, and its impacts and implications without management. A better understanding of the current situation of China's domestic marine fisheries, especially the little-understood component used as feed, can provide a solid basis for supporting future policy reforms in China's marine fisheries sector.

## 2 | METHODOLOGY

The present study adopted an integrated, interdisciplinary methodology by combining both top-down and bottom-up approaches to develop a holistic understanding of the feed component of China's domestic marine catch. We started from systematic literature reviews and then proceeded to a nationwide field survey of fishing vessels/gears and analysis of feed-grade fish samples, collected in eight major fishing provinces (Tables S2 and S3; Figure S3).

Field survey and feed-grade fish sampling were focused on the most important fishing months from August to December in 2016,

which is right after the national fishing moratorium and prior to the winter when fishing activities were largely closed in the north. In 2016, the total capture fisheries production (both marine and freshwater) in the second half of the year was 11.69 mmt, which accounts for 66.4% of annual production (Economic Daily, 2016; MOA, 2017a).

## 2.1 | Terms used

In the present study, domestic marine fisheries catches were classified into three categories: (a) *food species*; (b) *forage species*, particularly small pelagic and planktivorous species; and (c) all other *mixed species* (Table 1). In relation to supplying the animal feed sector, we suggest that the term “feed-grade fish” for catch was used as animal feed (either directly or indirectly after reduction), which consists of a wide range of species and sizes from small-sized individuals of food species (category a), forage species (category b) and mixed species (category c). Specifically, “feed-grade fish” refers to the mixture of poorly preserved, small-sized and low commercial-value fishes and invertebrates primarily used as animal feed. The term “feed-grade fish” refers to “Xiao Za Yu” in Mandarin, which literal translation is “small and miscellaneous fish”.

## 2.2 | Fishing vessel/gear survey

Fishing vessel/gear surveys were conducted in December 2016. The sampling of fishing vessels for interviews followed a multistage sampling approach from provincial level to district and port levels. Eight major marine fisheries provinces were selected for the survey: Liaoning, Shandong, Jiangsu, Zhejiang, Fujian, Guangdong, Guangxi and Hainan. The total marine catch in these eight provinces represented 97.7% of the national domestic marine catch in 2016 (Table S2). The most important one or two districts for marine capture fisheries in each province, in terms of the number of fishing vessels and landing volumes, giving a total of 22 fishing ports in 12 districts, were selected for the surveys (Table S3; Figure S3).

Surveys covered the following areas of interest: (a) basic information about interviewees such as position, name, age and contact

information; (b) vessel information such as registration number, length and horsepower; (c) major fishing areas and major fishing gears; (d) total catch of food species in different taxonomic groups (i.e., fishes, cephalopods, crustaceans and others); (e) total catch of feed-grade fish (i.e., the fish is marketed as such); (f) catches of major species among food species; (g) major species of feed-grade fish; and (h) destination and price of feed-grade fish. Survey questions were designed to cover the whole fishing season in the second half of 2016 (August–December).

Fishing vessel surveys at port level were primarily conducted according to fishing gear type. Fishing gears were classified according to The International Standard Statistical Classification of Fishing Gear (FAO, 2016) and standards in publications (Hainan Fisheries Institute, 2013; Sun, 2014; Wang & Yu, 2012). Survey sample sizes for each fishing gear category were at least 30, except for trawler. The trawler was the focus of the present study because trawlers produce half of China's marine catch by weight in the EEZ (Figure S4) and account for a high proportion of feed-grade fishes (Table S1). Survey sample sizes for trawler in each selected province were at least 30. The term “trawler” covers vessels that may target feed-grade fish (especially midwater trawlers) and those that catch feed-grade fish incidentally (generally bottom trawlers). Fishing vessels were randomly selected in each port, with interviewees being mainly fishing vessel owners, captains or senior fishers.

## 2.3 | Fish sampling, identification and measurement

From August to December 2016, feed-grade fish samples of up to 1 kg in weight were collected 2–3 times per month in each of eight provinces directly from fishing vessels. Species were identified, and individuals were counted and measured for body weight and length. Identified species from collected samples were classified as *food species*, *forage species* or *mixed species* based on the opinions of four experts from different research institutes. Classifications also referenced major aquatic trading websites, and marine capture species lists in the FAO FishstatJ database (FAO, 2017), China Fishery Statistical Yearbook (MOA, 2017a), the list of

**TABLE 1** Characteristics of three fisheries catch categories

Category	Food species	Forage species	Mixed species
Targeted	Yes	Yes	No
Utilization <sup>a</sup>	Human food	Mostly feed	Mostly feed
Production scale	Small to large	Small to large	Small to large
Commercially importance	High	Medium	Low
Planktivorous	Yes/no	Mostly yes	Mostly no
Pelagic/demersal	Demersal/pelagic	Mostly pelagic	Mostly demersal
Market value	Medium to high	Low to medium	Low
Primary fishing gear	All	Midwater trawl	Bottom trawl
Example species	Groupers, yellow croaker ( <i>Larimichthys polyactis</i> )	Anchovies, sardines	Pinkgray goby ( <i>Amblychaeturichthys hexanema</i> )

<sup>a</sup>Marketed and used as such, and dependent on size of fish, quality of handling, volume available, time of year and location.

commercially important aquatic animals and plants under special national protection for resources (MOA, 2017b), the standard of catchable size of important fisheries species part 1 (economic marine species; MOA, 2017c), and marine capture species information in FishBase (Froese & Pauly, 2017). The IUCN Red List Status of Threatened Species ([www.iucnredlist.org](http://www.iucnredlist.org)) was referred to for conservation status of identified species. Stock assessments from fisheries in China or the wider SE Asia region were collected from FAO evaluations and related publications.

## 2.4 | Data analysis

A multiple linear regression model was used to investigate factors contributing to different proportions of feed-grade fish in catches using SPSS 22 statistics software (IBM New York, NY, USA). All fishing vessels surveyed were classified into one of three length categories (<12 m, 12–24 m, and  $\geq 24$  m) according to the China Fishery Statistical Yearbook (MOA, 2017a). The proportion of feed-grade fish in catches was calculated based on survey results for all major fishing gears surveyed across all eight provinces. Using marine fisheries production by province and fishing gear type as a weighting factor for proportion of feed-grade fish, the overall proportion of feed-grade fish for the whole country was calculated. Independent-samples Kruskal–Wallis test was used for significance test using SPSS 22 statistic software. Values are given as means  $\pm$  SD and significance level was set at  $p < .05$ . The Chinese Yuan (CNY) was converted to US dollars (USD) using the annual average exchange rate in 2016 as 1 USD = 6.66 CNY (<https://www.investing.com/>).

Maturity stage was inferred from knowledge of recorded or estimated size (body length) at first maturity (Lm). Lm was collected from FishBase (Froese & Pauly, 2017). When the recorded Lm for a species was missing, the estimated Lm in the FishBase Species Ecology Matrix was used; estimated Lm was calculated from the recorded asymptotic length of a species and an empirical relationship between length at first maturity and asymptotic length (Froese & Binohlan, 2000; Froese & Pauly, 2017).

Several potential sources of variation and uncertainty in the present study (Appendix S1) could affect the calculations of total feed-grade fish volume as well as feed-grade fish sampling. To estimate the aggregate impact of data uncertainty on feed-grade fish production in China, a Monte Carlo analysis was conducted in Oracle Crystal Ball (Release 11.1.2.3.500). The Monte Carlo method is extensively used in fisheries research to estimate uncertainty by repeated random sampling based on predefined parameters of probability distributions (Pauly et al., 2014). The type of probability distributions and uncertainty parameters for the model were calculated by the Batch Fit module in Oracle Crystal Ball and based on survey data of the present study. The uncertainty parameters represented variations in the proportion of feed-grade fish from different fishing gear categories. We calculated uncertainty based on a distribution of 100,000 runs resulting from the Monte Carlo sampling method.

A sensitivity analysis was conducted to explore the juvenile percentages associated with variation of length at sexual maturity (Lm)

(Appendix S1). Instead of using the estimated Lm, body length at 50% of the maximum body length (L50) reported in the FishBase (Froese & Pauly, 2017) was used as a comparable criterion for the size of sexual maturation.

## 3 | RESULTS

### 3.1 | Proportion of feed-grade fish in catch

A total of 886 interviews were conducted from domestic vessels in 22 major fishing ports in eight major fishing provinces in December 2016 (Tables S3 and S4). The multiple linear regression model shows fishing gear and sampling location (province) to be the major factors associated with different proportions of feed-grade fish (including fishes and invertebrates) in catches ( $p < .05$ ; Table S5). The average engine power of fishing vessels, according to respondents willing and able to answer this question, was  $410.5 \pm 343.3$  kW ( $n = 582$ ), and the average vessel length is  $31.3 \pm 10.0$  m ( $n = 851$ ; Figure S5). Although large fishing vessels produced significantly more catch per vessel than small fishing vessels (Table S6), vessel size (length) has little effect on the proportion of feed-grade fish in catches (Table S7).


Trawls, stow nets, gillnets and entangling nets, and surrounding nets and seine nets were identified as the main fishing gears taking feed-grade fish. Trawls accounted for the highest proportion of feed-grade fish catch overall by weight (Figure 1), with nearly half ( $48.9 \pm 26.5\%$ ,  $n = 538$ ) of trawler catches being feed-grade fish. Midwater trawls primarily target small pelagic forage fish species, mainly anchovies in the Bohai and Yellow Seas, producing  $71.5 \pm 31.2\%$  ( $n = 76$ ) of feed-grade fish as a proportion of their total catch by weight. This was significantly ( $p < .05$ ) higher than the proportion taken from bottom trawlers ( $43.6 \pm 21.3\%$ ,  $n = 336$ ; Table 2).

Three-fifths (58.5%) of trawlers, 46.6% of stow net vessels, 23.5% of gillnet and entangling net vessels, and 7.1% surrounding nets and seine nets vessels by number were predominantly producing feed-grade fish, accounting for at least 50% of their total catch by weight (Figure 2). Trawlers in Liaoning, Shandong, Guangxi and Hainan reported significantly higher proportions of feed-grade fish in their catches than other provinces ( $p < .05$ ; Table 3).

### 3.2 | Feed-grade fish production

The overall proportions of feed-grade fish, according to 886 interviews, weighted by production by gear and province (Tables 4 and 5) show that 34.7% of China's marine catch, or 4.6 mmt marine fisheries production, was feed-grade fish. The Monte Carlo analysis also shows that about 4.6 mmt (95% confidence interval: 2.1–7.0 mmt) of total production was feed-grade fish (Figure S6, Table S8).

The proportion of feed-grade fish produced by all trawlers combined, weighted by production by province, was 48.7%, which is equivalent to 3.0 mmt of feed-grade fish production from China's EEZ (Table 5). This is followed by stow nets 0.7 mmt, gillnets and entangling nets 0.6 mmt, and surrounding nets and seine nets 0.2 mmt (Table 5). Shandong, Guangxi and Zhejiang had the highest



**FIGURE 1** Proportions of feed-grade fish in catches (by weight) from different fishing vessels classified by fishing gears and transport ship. “Other” category includes hooks and lines, pots, rakes and other miscellaneous gears

proportions of feed-grade fish (Figure 3) in their catches, reflecting the reliance of these provinces on trawlers (Figure S7). The top capture fisheries provinces are also major feed-grade fish-producing provinces, especially Zhejiang and Shandong (Figure 4; Table S2).

### 3.3 | Use and prices of feed-grade fish

A total of 88.2% of interviewees ( $n = 781$ ) reported that their feed-grade fish catch was mainly used in the aquaculture and feed industries, either as direct feed (e.g., using unprocessed whole fish or fish parts directly in aquaculture cages/ponds) or indirectly to produce fishmeal and fish oil which is then used as raw material for aquaculture and livestock feed (Figure 5). Some mink and fox farms in Shandong and Liaoning, and, occasionally, chicken and pig farms in fish-producing areas, also use feed-grade fish but the scale of this use is not known. A low proportion (4.9%)

of the interviewees ( $n = 43$ ) reported discarding feed-grade fish because of poor condition, while 6.7% ( $n = 59$ ) reported their catch was mainly used for human consumption after processing into minced products such as fish balls, dried or salted product, or sauces and pastes. The consumption of feed-grade fish (4.6 mmt) was estimated according to survey data (Figure 6).

Prices of feed-grade fish purchased from fishing vessels or at fishing ports varied from USD 0.1–0.8 per kg and averaged USD  $0.2 \pm 0.1$  per kg ( $n = 684$ ) depending on province, freshness/quality, species composition, the type of utilization intended, and the demand for feed-grade fish by the aquaculture sector (Figure S8). Small size and poor condition of feed-grade fish led to particularly low prices, making them affordable for purchase and use whole by aquaculture producers or for manufacturing of fishmeal.

A nationwide feed-grade fish value chain has developed, and a stable market and price structure for feed-grade fish now exists. Feed-grade fish transactions take place both within the capture fisheries sector and across a trade chain that links feed-grade fish capture to transportation networks by sea or by land and involving cold storage, with aquaculture farms and fishmeal factories as major end-users. The trade network runs across the country; for example, feed-grade fish caught in Zhejiang could supply aquaculture farms in Fujian, and fishmeal factories in Shandong.

**TABLE 2** Proportions of feed-grade fish in catches (by weight) from different fishing gears (mean  $\pm$  SD,  $n = 886$  interviews)

Vessel type	Proportion of feed-grade fish (%)
Midwater trawler	$71.51 \pm 31.20$ ( $n = 76$ ) <sup>a</sup>
Bottom trawler	$43.59 \pm 21.25$ ( $n = 336$ ) <sup>b</sup>
Unspecified trawler <sup>†</sup>	$49.57 \pm 29.01$ ( $n = 126$ ) <sup>b</sup>
Stow nets	$40.94 \pm 26.31$ ( $n = 151$ ) <sup>b</sup>
Gillnets and entangling nets	$27.54 \pm 25.26$ ( $n = 55$ ) <sup>cd</sup>
Surrounding nets and seine nets	$17.18 \pm 19.21$ ( $n = 87$ ) <sup>d</sup>
Other <sup>‡</sup>	$7.63 \pm 6.37$ ( $n = 19$ ) <sup>d</sup>
Transport ship	$43.50 \pm 29.56$ ( $n = 36$ ) <sup>bc</sup>

Note: Means superscripted by different letters indicate significant differences ( $p < .05$ ). Independent-samples Kruskal–Wallis test was used for significance test using SPSS 22 statistic software.

<sup>†</sup>Unspecified trawler includes both midwater trawler and bottom trawler, but not specified by the interviewees.


<sup>‡</sup>“Other” category includes hooks and lines, pots, rakes and other miscellaneous gears.

### 3.4 | Species composition

A total of 80 samples of feed-grade fish, with a total weight of 64.4 kg, after removing unidentifiable fish and invertebrate body parts and garbage, were collected from 22 fishing ports surveyed throughout eight major marine fisheries provinces. A total of 12,206 individuals were measured, the majority being fish, with 10,796 (88.5%) individuals, followed by 1,263 (10.4%) crustaceans, and 147 (1.2%) cephalopods.

Species identification focused on fishes; 98.0% of fish individuals were identified to species level. A total of 218 fish species were identified (three to genus level only), including three Chondrichthyans (two orders and two families) and 215 Actinopterygians (19 orders and 84

**FIGURE 2** Proportions of feed-grade fish in catches (by weight) from all types of trawlers in eight provinces (a) by proportion of feed-grade fish and (b) by province ( $n = 538$ )



families) (Figure 7; Figure S9, Tables S9–S11). The most numerous fish families were Engraulidae (3,668 individuals from 10 species), Gobiidae (1,852 individuals from 14 species), Clupeidae (752 individuals from eight species), Synodontidae (734 individuals from five species), Bregmacerotidae (632 individuals from one species) and Sciaenidae (488 individuals from 17 species). In addition, 50 crustaceans (two orders and 20 families) and five cephalopods were identified (Tables S12 and S13). Many crustaceans and cephalopods were small-sized and/or body damaged; therefore, body size/weight measurements were not made.

### 3.5 | Food fish species and proportion of catches composed of juveniles

Of 218 identified fish species, 102 (46.8%) were food species, 19 (8.7%) were forage fish and 97 (44.5%) mixed fish. Of the 10,580 fish individuals identified to species, 3,112 (29.4%) were food fish, 4,392 (41.5%) forage fish, and 3,076 (29.1%) mixed fish. A total of 9,377 (88.6%) individuals had measurable body length. Three-quarters (7,225 individuals, 77.05%) of these fish individuals were estimated to be in the juvenile phase in terms of the criterion of  $L_m$  (Tables

S9–S11). The majority (88.9%) of food fish were likely to be in their juvenile stage while 71.8% and 73.1% of forage fish and mixed fish were juveniles, respectively (Figure 8).

The percentages of juveniles were lower (food fish 79.7%, forage fish 60.9%, mixed fish 37.3% and overall 59.9%) when the criterion for size of maturity was set for the criterion of  $L_{50}$  (Figure S10). The reduced percentage of juveniles is mainly because the estimated  $L_m$  tends to be longer in small species than in large species (Froese & Binohlan, 2000; Froese & Pauly, 2017; Tsikliras & Stergiou, 2014). Most of the fish species identified in this study are small-sized species (Tables S9–S11), and their estimated  $L_m$ s are more likely longer than the sizes of 50% of the maximum body length.

### 3.6 | Stock assessment and conservation status

Of 218 fish species identified, stock assessments published in reports, books or papers from 2005 to 2015 were found for 32 species and taxonomic groups (to genus level), of which 28 (87.5%) were over-exploited, and 4 (12.5%) were fully exploited or starting to decline at the time of reporting, all were Actinopterygians (Tables S14 and S15).

**TABLE 3** Proportions of feed-grade fish in catches (by weight) from trawlers in eight provinces (mean  $\pm$  SD,  $n = 538$  interviews)

Province	Bottom trawler (%)	Midwater trawler (%)	Unspecified trawler (%) <sup>†</sup>
Liaoning	36 $\pm$ 35 ( $n = 15$ ) <sup>b</sup>	94 $\pm$ 8 ( $n = 21$ ) <sup>a</sup>	23 $\pm$ 40 ( $n = 3$ ) <sup>abc</sup>
Shandong	35 $\pm$ 26 ( $n = 41$ ) <sup>b</sup>	85 $\pm$ 13 ( $n = 27$ ) <sup>ab</sup>	67 $\pm$ 23 ( $n = 15$ ) <sup>a</sup>
Jiangsu	33 $\pm$ 29 ( $n = 15$ ) <sup>b</sup>		33 $\pm$ 33 ( $n = 9$ ) <sup>abc</sup>
Zhejiang	56 $\pm$ 13 ( $n = 51$ ) <sup>a</sup>	57 $\pm$ 16 ( $n = 5$ ) <sup>bc</sup>	35 $\pm$ 32 ( $n = 24$ ) <sup>bc</sup>
Fujian	40 $\pm$ 18 ( $n = 63$ ) <sup>b</sup>	26 $\pm$ 34 ( $n = 14$ ) <sup>c</sup>	9 $\pm$ 12 ( $n = 10$ ) <sup>c</sup>
Guangdong	31 $\pm$ 13 ( $n = 73$ ) <sup>b</sup>		0 $\pm$ 0 ( $n = 3$ ) <sup>bc</sup>
Guangxi	58 $\pm$ 11 ( $n = 63$ ) <sup>a</sup>	51 $\pm$ 0 ( $n = 3$ ) <sup>bc</sup>	60 $\pm$ 9 ( $n = 11$ ) <sup>ab</sup>
Hainan	62 $\pm$ 11 ( $n = 15$ ) <sup>a</sup>	62 $\pm$ 27 ( $n = 6$ ) <sup>bc</sup>	65 $\pm$ 14 ( $n = 51$ ) <sup>a</sup>

Note: Means superscripted by different letters indicate significant differences ( $p < .05$ ). Independent-samples Kruskal–Wallis test was used for significance test using SPSS 22 statistic software.

<sup>†</sup>Unspecified trawler includes both midwater trawler and bottom trawler but not specified by the interviewees.

In terms of conservation status, 143 fish species have not been evaluated (NE) against the IUCN Red List Criteria. Among the 72 species evaluated, one was endangered (EN, Threadfin porgy [*Evynnis cardinalis*]), one vulnerable (VU, Golden threadfin bream [*Nemipterus virgatus*]), 59 least concern (LC) and 11 data deficient (DD; Tables S9–S11).

Eight species of fish, one crustacean and one cephalopod have been important for China's national stock enhancement projects (Table S16). Of these ten species, most individuals taken in samples of the current study were still in their juvenile stage. National stock enhancement projects, therefore, are likely being compromised by uncontrolled fisheries that catch feed-grade fish as these projects rely on the dispersal of juvenile fish into the wild.

## 4 | DISCUSSION

### 4.1 | Fishing for feed in China is substantial and biologically unsustainable

Although the reported production from China's fishing grounds remained stable over multiple years, an increasing and significant proportion of domestic marine catch comprises poorer quality fish, less marketable as direct human food than previously, and now primarily used as animal feed. The loss of historically major food fisheries, declining fisheries biomass and CPUE, changing species composition, declining fish size, and shifting end use of marine catch, primarily driven by severe overfishing and excessive capture of juveniles,

are trends that have largely been obscured by this stable reported production.

At the global level, around one-quarter, or 20 mmt, of fisheries landings were used for purposes other than direct human consumption (FAO, 2018). The present study found that about 35% (4.6 mmt) of China's marine catch is now sold at low prices as feed-grade fish, 40% higher than the global average. Considering China's feed-grade fish production alone would rank it fourth among world's leading marine fisheries countries (Figure 9).

Thirty per cent (29.4%) of feed-grade fish catch by number of individuals was identified as food fish species, with most of these (88.9%) in their juvenile size range. If allowed to reach marketable size, these fish could potentially have gone directly to human food use and gained higher prices, which in turn would maximize yield or, at least, optimize individual value per recruit (Beverton & Holt, 1993; Cashion, Le Manach, Zeller, & Pauly, 2017; Froese, Winker, Gascuel, Sumaila, & Pauly, 2016; also see Table S17). Forage fish are also overfished in China, with 71.8% of the individuals sampled being juveniles. This is in contrast to many of the world's major reduction fisheries that specifically target forage fish which are effectively managed using a variety of catch control tools, such as quotas and effort limits (Pikitch et al., 2012; Tveteras, Paredes, & Peña-Torres, 2011).

Protecting juveniles is a core component of fisheries management and sustainable resource use (Crowder et al., 2008; Froese et al., 2016; Garcia, Zerbi, Aliaume, Do Chi, & Lasserre, 2003; Pinsky, Jensen, Ricard, & Palumbi, 2011). There are examples of fisheries collapses of multiple different individual species stocks associated with depletion of adult individuals and an increasingly high proportion of juveniles in catches. For example, the large yellow croaker (*Larimichthys crocea*) was heavily exploited to the point of severe depletion in the 1980s with an increasingly heavy take of juveniles (Liu & Sadovy de Mitcheson, 2008; Figure S11). Trends of increasingly higher proportions of juveniles in catches were also observed with many other major food species such as largehead hairtail (*Trichiurus lepturus*), yellow croaker (*Larimichthys polyactis*), silver pomfret (*Pampus argenteus*) and Japanese Spanish mackerel (*Scomberomorus niphonius*; Figure S12).

The outcomes of decades of overfishing and excess fishing capacity in China's domestic marine fisheries resulted in depletions of some "traditionally taken" species by the 1970s and 1980s. By the 1980s and 1990s, some forage species and recently exploited food species such as the scaly hairfin anchovy (*Setipinna taty*) and the filefish (*Thamnaconus modestus*) had also become over-exploited (Figure S2). The mean trophic level of China's marine catch has declined from 3.70 to 3.46 during the past half-century (Cao et al., 2017), which is linked to the expanding proportion of low-trophic-level forage species in catches, eroding biodiversity and degrading marine ecosystem structure, functioning and resilience (Smith et al., 2011). Unmanaged over the long term, there is a real risk that the fishery resources underpinning the feed component of China's marine catch will also inevitably collapse (Beverton, 1990; Costello, 2017; Figure S2).



**TABLE 4** Weighting factors and weighted proportion of feed-grade fish in different fishing gears

Fishing gear	Province	Proportion of feed-grade fish collected from interview (Pp, %)	Marine catch in China's EEZ in 2016 (Pd, mt) <sup>a</sup>	Weighting factors (Wt) <sup>b</sup>	Weighted proportion of feed-grade fish of each fishing gear category (WPP, %) <sup>c</sup>
Trawls	Liaoning	66.01	397,894	0.065075	48.70
	Shandong	56.91	1,398,197	0.228673	
	Jiangsu	32.85	78,717	0.012874	
	Zhejiang	49.62	2,103,225	0.343980	
	Fujian	33.80	778,343	0.127297	
	Guangdong	29.70	749,572	0.122592	
	Guangxi	58.20	425,929	0.069660	
	Hainan	63.76	182,511	0.029849	
Surrounding nets and seine nets	Shandong	60.01	36,735	0.033672	20.34
	Jiangsu	55.56	5,724	0.005247	
	Zhejiang	15.00	220,102	0.201749	
	Fujian	1.51	311,547	0.285569	
	Guangdong	37.50	143,089	0.131158	
	Guangxi	25.18	62,174	0.056990	
	Hainan	28.77	311,597	0.285615	
Gillnet	Liaoning	8.33	491,656	0.220880	20.20
	Shandong	39.24	416,024	0.186902	
	Jiangsu	35.26	163,291	0.073360	
	Zhejiang	27.88	319,057	0.143339	
	Fujian	2.00	318,118	0.142917	
	Guangdong	3.89	433,718	0.194851	
	Guangxi	90.00	84,030	0.037751	
Stow nets	Liaoning	21.92	95,891	0.065429	47.53
	Shandong	51.69	190,367	0.129892	
	Jiangsu	31.00	211,978	0.144638	
	Zhejiang	47.18	613,130	0.418355	
	Fujian	62.82	346,156	0.236191	
	Guangdong	58.64	8,052	0.005494	
Others <sup>d</sup>	All	0.00	1,360,431	—	0

<sup>a</sup>Data source: (MOA, 2017a).

<sup>b</sup>Weighting factors ( $Wt_i$ ) are calculated separately for each fishing gear category and based on production in each province ( $Pd_i$ ), as  $Wt_i = Pd_i / \sum_{i=1}^n Pd_i$ , for each fishing gear category  $\sum_{i=1}^n Wt_i = 1$ .

<sup>c</sup>Weighted proportion of feed-grade fish of each fishing gear category (WPP) are calculated separately for each fishing gear category and based on proportion of feed-grade fish collected from interview ( $Pp_i$ ) and its weighting factors ( $Wt_i$ ), as  $WPP = \sum_{i=1}^n Pp_i \times Wt_i$ .

<sup>d</sup>"Others" category includes hooks and lines, pots, rakes and other miscellaneous gears. The proportion of feed-grade fish from "Others" category was set at 0% due to the limited sample size of it.

## 4.2 | Effective management is needed for both food and feed components

China has regulations, policies and measures aimed at controlling fishing capacity, including a fishing licence system, fishing moratorium and minimum mesh size regulation (Cao et al., 2017). However, these efforts have clearly been inadequate in part due to ineffective compliance/enforcement of regulations but also because these

measures do not control fishing effort. For example, along with depletion of traditional fisheries resources, fishers tend to reduce their mesh size to maintain or increase their catches (Yu, 2003); the legal minimum stretched mesh size requirement of 25 mm (which is still lower than 40 mm recommended by the APFIC Trawl Guidelines; FAO, 2014a) is poorly implemented. Cod-end mesh sizes net of 10 mm or less are extensively used by Chinese trawlers (Liang & Pauly, 2017).

**TABLE 5** Weighted means of proportion and production volume of feed-grade fish in total landings in China

Fishing gear	Weighted proportion of feed-grade fish (WPP, %) <sup>a</sup>	Marine catch of each fishing gear category in China's EEZ in 2016 (GPd, mmt) <sup>b</sup>	Overall weighting factors (OWt) <sup>c</sup>	Overall weighted proportion of feed-grade fish (OWPp, %) <sup>d</sup>	Feed-grade fish production (FFPd, mmt) <sup>e</sup>
Trawls	48.70	6.23	0.469	<b>34.69</b>	3.03
Surrounding nets and seine nets	20.34	1.10	0.083		0.22
Gillnets and entangling nets	20.20	3.04	0.229		0.61
Stow nets	47.53	1.55	0.117		0.74
Others <sup>f</sup>	0.00	1.36	0.102		0.00
Total catch in China's EEZ in 2016 (mmt)					13.28
Total feed-grade fish production in China in 2016 (mmt)					4.61

<sup>a</sup>The weighted proportion of feed-grade fish of each fishing gear category (WPP), please see Table 4.

<sup>b</sup>Data source: (MOA, 2017a).

<sup>c</sup>Overall weighting factors (OWt<sub>j</sub>) are calculated based on marine catch production of each fishing gear category in China's EEZ in 2016 (GPd<sub>j</sub>), as  $OWt_j = GPd_j / \sum_{j=1} GPd_j$ , and  $\sum_{j=1} OWt_j = 1$ .

<sup>d</sup>Overall weighted proportion of feed-grade fish (OWPp) are calculated based on weighted proportion of feed-grade fish of each fishing gear category (WPP<sub>j</sub>) and overall weighting factors (OWt<sub>j</sub>), as  $OWPp = \sum_{j=1} WPP_j \times OWt_j$ .

<sup>e</sup>Feed-grade fish production (FFPd) is calculated based on weighted proportion of feed-grade fish of each fishing gear category (WPP<sub>j</sub>) and marine catch production of each fishing gear category in China's EEZ in 2016 (GPd<sub>j</sub>), as  $FFPd = WPP_j \times GPd_j$ .


<sup>f</sup>"Others" category includes hooks and lines, pots, rakes and other miscellaneous gears. The proportion of feed-grade fish from "Others" category was set at 0% due to the limited sample size of it.


China's fisheries resources have been over-exploited since the 1970s, well before the development of intensive feed and aquaculture industries. Indeed, the aquaculture industry could be considered a beneficiary of changed marine catch composition. Although aquaculture development and intensification was not the direct cause of over-exploited fisheries resources in China, the rapid development of the feed and aquaculture industries in China fuelled a high market demand for feed-grade fish and provided new incentives to continue fishing even as traditional fisheries were being depleted.

With globally expanding aquaculture, fishmeal prices in the international market have more than tripled over the past decade

(FAO, 2014b; also see Figure S13), and the price of domestically produced fishmeal has increased in tandem. The high demand for fish feed from the aquaculture sector is supported by a large number of fishmeal factories. Their operation, often well below full capacity (20%–45% in 2016; Wang, 2017), suggests increasing competition for scarce supplies. Such pressures place further stress on over-exploited fishery resources.

The fishery fuel subsidy policy was introduced in 2006 in China and became the most important type of fishery subsidy in the country, accounting for 94% of overall fishery subsidies (Mallory, 2016). The fuel subsidy covers about one-third of the total cost of marine capture fisheries (Zhu & Huang, 2014). Without fuel subsidies, more

**FIGURE 3** Proportions of feed-grade fish in marine fisheries catches in different provinces





**FIGURE 4** Contribution of feed-grade fish production (by weight proportion) in China by major producing province


than 90% of fishing vessels would lose money (Guo, 2015). Economic incentives, from increasing demand for feed-grade fish, high fuel subsidies, along with depletion of traditional fishing target species, further increase the motivation for overfishing of both food and feed species.

The unparalleled scale (i.e., high volume) and poor fisheries resources condition (i.e., highly depleted) of China's domestic marine fisheries make them distinctive and the solutions to the problems created need to be large scale and must significantly reduce current levels of fishing effort. China, like many other tropical and subtropical Asian countries (Leadbitter, 2013), has a high biodiversity of fishes. The best option for managing the take of hundreds of different species, on the one hand, while addressing conservation principles that reduce threats to species and maintain functional marine ecosystems, may not be to seek Maximum Sustainable Yield (see for example Sainsbury, 1984). One alternative is to generate a "Pretty Good Yield" (Rindorf, Cardinale, et al., 2017; Rindorf, Dichmont, et

**FIGURE 5** Feed-grade fish consumption and other use, in different provinces. Use of feed-grade fish as direct feed or to produce fishmeal was not specified in surveys conducted in Zhejiang province




**FIGURE 6** Estimated production and consumption of marine catch in China's EEZ. The proportions of forage and mixed species are based on data in Figure S2; the use of feed-grade fish in Zhejiang was set at national average (Figure 5); the unit for all numbers is mmt. Figure appears in colour in the online version only




**FIGURE 7** Taxonomic coverage of feed-grade fish samples. (a) Class, (b) order and (c) family

**FIGURE 8** Number and the proportion of juveniles and adults in the samples in three categories (food fish, forage fish and mixed fish). Unidentified individuals are those individuals identified to species level, but no available measurable body length



**FIGURE 9** Production of top 10 world's leading marine (domestic + distant) fisheries countries and production of feed-grade fish in China's marine catch in 2016. Data source: (FAO, 2017); aquatic plants excluded



al., 2017) condition with acceptable trade-offs where marine ecosystems are able to satisfy diverse societal needs ensuring most stocks and the ecosystem are protected or remain biologically viable (Cheung & Sumaila, 2008; Costello, 2017; FAO, 2014a). Conservation measures, such as marine protected areas, can be used to safeguard key spawning and nursery areas that enable species to replenish themselves (Edgar et al., 2014; Hastings, Gaines, & Costello, 2017; Roberts et al., 2017).

By reducing excess fishing capacity, safeguarding key life history times and/or places, and effectively enforcing the current minimal mesh size requirement, fishery collapse could be averted; fisheries biomass, total catch and value could be increased; and the catch profile could be shifted to favour a greater proportion of larger fish (Costello, 2017; Jacobsen, Gislason, & Andersen, 2013; Szuwalski, Burgess, Costello, & Gaines, 2017). This could also produce a better mix of products for direct human consumption, and indirectly via fish and livestock feed.

## 5 | CONCLUSIONS

Significant changes in species composition in catches, the high proportion of catch comprised of juveniles, and loss of or severe declines in traditionally important fisheries are clear evidence of weak fisheries management and a decades-long history of persistent overfishing and excess fishing capacity in China's domestic marine fisheries. The growing proportion of catch used to supply the aquaculture sector since the 1990s has meant that pressure continues to be high across the coastal ecosystem, even as these declines and shifts occur. To date, commitments to resolve fishery issues remain unfulfilled and existing management measures have been ineffective.

Under current fishery policy, China's fisheries risk a further downwards shift due to intensification of both fishing down and farming up the food web (Naylor et al., 2000; Pauly et al., 1998; Tacon, Metian, Turchini, & De Silva, 2010). Unchecked, this

trajectory will continue and could ultimately result in an ecosystem shift towards species that have little or no human value and other challenges of ecosystem degradation, some of which may not be reversible, and which may not even be able to supply feed-grade fish into the future. There is also a long-term threat associated with loss of marine biodiversity and loss of ecosystem function and resilience.

Based on the outcomes of the present study, it is clear that the feed component of China's marine catch should be documented and managed as part of the overall national planning for sustainable fisheries resources use and marine ecosystem management. A biologically sustainable balance between fishing for food and fishing for feed could be achieved to produce a better mix of products by effectively managing China's multispecies fisheries and reducing excessive fishing capacity, while not further compromising the ecosystem and associated biodiversity. Overall, a more ecosystem-level and precautionary approach is needed for fisheries management, aquaculture development (such as a better selection of species cultured), including management of the feed component of China's marine catch to ensure long-term sustainability.

## ACKNOWLEDGEMENTS

This study was motivated by the two symposiums on "Fisheries and Food Security in China" hosted by Stanford University's Center on Food Security and the Environment (FSE). Wenbo Zhang is supported by the China-ASEAN Maritime Cooperation Fund. Min Liu is supported by the National Natural Science Foundation of China (No. 41476149) and the State Oceanic Administration of China (220203993022761133). Yvonne Sadovy de Mitcheson received funds from ADM Capital Foundation and Swire Trust. Thanks to Zhuocheng Zhou, Ruiqiang Zheng and Xiaoyun Tang for field survey and specimen collection, to more than 20 volunteers and employees in local fisheries cooperatives, and to Weidi Yang and postgraduates at Fish Biology Laboratory of Xiamen University for species identification and measurement assistant. Comments and suggestions for the original research report from anonymous experts at the Chinese Academy of Fishery Sciences, Shanghai Ocean University, and Xiamen University are much appreciated. Figure 6 produced using Sankey Flow Show software (<https://www.sankeyflowshow.com/index.html>).

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ORCID

Wenbo Zhang  <https://orcid.org/0000-0003-3765-3980>

Min Liu  <https://orcid.org/0000-0002-6768-6794>

Yvonne Sadovy de Mitcheson  <https://orcid.org/0000-0002-1154-3081>

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## REFERENCES

- Alverson, D. L., Freeberg, M. H., Murawski, S. A., & Pope, J. G. (1994). *A global assessment of fisheries bycatch and discards (FAO Fisheries Technical Paper No. 339)*. Retrieved from <http://www.fao.org/docrep/003/T4890E/T4890E00.HTM>
- Anon (2016). *Fisheries Statistics of Thailand 2014. No. 11/2016, Information and Communication Technology Center, Department of Fisheries, Ministry of Agriculture and Cooperatives*.
- Beverton, R. J. H. (1990). Small marine pelagic fish and the threat of fishing; are they endangered? *Journal of Fish Biology*, 37, 5–16. <https://doi.org/10.1111/j.1095-8649.1990.tb05015.x>
- Beverton, R. J. H., & Holt, S. J. (1993). *On the dynamics of exploited fish populations* (1st. ed.). Dordrecht: Springer. <https://doi.org/10.1007/978-94-011-2106-4>
- Cao, L., Chen, Y., Dong, S., Hanson, A., Huang, B. O., Leadbitter, D., ... Naylor, R. L. (2017). Opportunity for marine fisheries reform in China. *Proceedings of the National Academy of Sciences*, 114(3), 435–442. <https://doi.org/10.1073/pnas.1616583114>
- Cao, L., Naylor, R., Henriksson, P., Leadbitter, D., Metian, M., Troell, M., & Zhang, W. (2015). China's aquaculture and the world's wild fisheries. *Science*, 347(6218), 133–135. <https://doi.org/10.1126/science.1260149>
- Cashion, T., Le Manach, F., Zeller, D., & Pauly, D. (2017). Most fish destined for fishmeal production are food-grade fish. *Fish and Fisheries*, 18(5), 837–844. <https://doi.org/10.1111/faf.12209>
- Chau, G. T. H., & Sadovy, Y. (2005). The use of mixed fish feed in Hong Kong's mariculture industry. *World Aquaculture*, 36(4), 6.
- Chee, P. E. (1997). A review of the bycatch and discards in the fisheries of Southeast Asia. In I. J. Clucas, & D. G. James (Eds.), *Papers presented at the technical consultation on reduction of wastage in fisheries*. Tokyo. *FAO Fisheries Report No. 547 (Suppl.)* (pp. 257–274). Tokyo, Japan: Rome, FAO.
- Cheng, J., & Yu, L. (2004). The change of structure and diversity of demersal fish communities in the Yellow Sea and East China Sea in winter (in Chinese). *Journal of Fisheries of China*, 28(1), 29–34. <https://doi.org/10.3321/j.issn:1000-0615.2004.01.005>
- Cheung, W. W. L., & Sadovy, Y. (2004). Retrospective evaluation of data-limited fisheries: A case from Hong Kong. *Reviews in Fish Biology and Fisheries*, 14(2), 181–206. <https://doi.org/10.1007/s11160-004-5422-y>
- Cheung, W. W. L., & Sumaila, U. R. (2008). Trade-offs between conservation and socio-economic objectives in managing a tropical marine ecosystem. *Ecological Economics*, 66(1), 193–210. <https://doi.org/10.1016/j.ecolecon.2007.09.001>
- Chiu, A., Li, L., Guo, S., Bai, J., Fedor, C., Naylor, L. R., & Naylor, R. L. (2013). Feed and fishmeal use in the production of carp and tilapia in China. *Aquaculture*, 414–415, 127–134. <https://doi.org/10.1016/j.aquaculture.2013.07.049>
- Costello, C. (2017). Fish harder; catch more? *Proceedings of the National Academy of Sciences*, 114(7), 1442–1444. <https://doi.org/10.1073/pnas.1620731114>
- Crowder, L. B., Hazen, E. L., Avissar, N., Bjorkland, R., Latanich, C., & Ogburn, M. B. (2008). The impacts of fisheries on marine ecosystems and the transition to ecosystem-based management. *Annual Review of Ecology, Evolution, and Systematics*, 39(1), 259–278. <https://doi.org/10.1146/annurev.ecolsys.39.110707.173406>
- Economic Daily (2016). The main indicators of fishery in China have improved in the first half of 2016 (in Chinese). *Rural Know-All*, (20), 13.

- Edgar, G. J., Stuart-Smith, R. D., Willis, T. J., Kininmonth, S., Baker, S. C., Banks, S., ... Thomson, R. J. (2014). Global conservation outcomes depend on marine protected areas with five key features. *Nature*, 506(7487), 216–220. <https://doi.org/10.1038/nature13022>
- Edwards, P. (2008a). Rural aquaculture: from integrated carp polyculture to intensive monoculture in the Pearl River Delta, South China. *Aquaculture Asia Magazine*, (June), 3–7. Retrieved from <http://library.enaca.org/AquacultureAsia/Articles/april-june-2008/1-peter-edwards-april-08.pdf>
- Edwards, P. (2008b). The changing face of pond aquaculture in China. *Global Aquaculture Advocate*, (September/October), 77–80. Retrieved from <http://pdf.gaalliance.org/pdf/GAA-Edwards-Sept08.pdf>
- Edwards, P., Tuan, L. A., & Allan, G. L. (2004). A survey of marine trash fish and fish meal as aquaculture feed ingredients in Vietnam. ACIAR Working Paper No. 57. Retrieved from <http://leanhtuan.vnweblogs.com/gallery/23458/03-trashfishreport-vietnamaciar.pdf>
- FAO (2007). *The state of world fisheries and aquaculture 2006*. Retrieved from <http://www.fao.org/docrep/009/A0699e/A0699e00.htm>
- FAO (2011a). Aquaculture development. 5. Use of wild fish as feed in aquaculture. In FAO. *Technical guidelines for responsible fisheries*. Retrieved from <http://www.fao.org/docrep/field/003/ab825f/AB825F00.htm#TOC>
- FAO (2011b). International Guidelines on bycatch management and reduction of discards. In FAO *international guidelines*. Retrieved from <http://www.fao.org/3/a-ba0022t.pdf>
- FAO (2014a). APFIC/FAO Regional Expert Workshop on “Regional guidelines for the management of tropical trawl fisheries in Asia”. Phuket, Thailand, 30 September–4 October 2013. In *RAP Publication 2014/01*. Retrieved from <http://www.fao.org/publications/card/en/c/f64766d6-7097-48cf-84fc-8fab66efb807/>
- FAO (2014b). *The state of world fisheries and aquaculture 2014*. Retrieved from <http://www.fao.org/3/a-i3720e/index.html>
- FAO (2016). *Revised international standard classification of fishing gears (ISSCFG, Rev.1 21 October 2010)* (p. 2). Rome, Italy: FAO.
- FAO (2017). *FishstatJ*. Retrieved from <http://www.fao.org/fishery/statistics/software/fishstatj/en>
- FAO (2018). *The state of world fisheries and aquaculture 2018 – Meeting the sustainable development goals*. 978-92-5-130562-1
- Froese, R., & Binohlan, C. (2000). Empirical relationships to estimate asymptotic length, length at first maturity and length at maximum yield per recruit in fishes, with a simple method to evaluate length frequency data. *Journal of Fish Biology*, 56(4), 758–773. <https://doi.org/10.1006/jfbi.1999.1194>
- Froese, R., & Pauly, D. (2017). *FishBase*. Retrieved from [www.fishbase.org](http://www.fishbase.org)
- Froese, R., Winker, H., Gascuel, D., Sumaila, U. R., & Pauly, D. (2016). Minimizing the impact of fishing. *Fish and Fisheries*, 17(3), 785–802. <https://doi.org/10.1111/faf.12146>
- Funge-Smith, S., Briggs, M., & Miao, W. (2012). Regional overview of fisheries and aquaculture in Asia and the Pacific 2012. In *RAP publication*. Retrieved from <http://www.fao.org/documents/card/en/c/951b0503-aece-5bdf-a4fa-500b868b55a1/>
- Funge-Smith, S., Lindebo, E., & Staples, D. (2005). Asian fisheries today: The production and use of low value/trash fish from marine fisheries in the Asia-Pacific region. In *Rap Publication, 2005/16*, 47.
- Garcia, S. M., Zerbi, A., Aliaume, C., Do Chi, T., & Lasserre, G. (2003). The ecosystem approach to fisheries. Issues, terminology, principles, institutional foundations, implementation and outlook. In *FAO Fisheries Technical Paper*. No. 443. <https://doi.org/10.1111/j.1467-2979.2010.00358.x>
- Guo, X. (2015). *Research on diesel subsidy policy influence on Ningbo ocean fishing (in Chinese)*. Shanghai, China: Shanghai Ocean University.
- Hainan Fisheries Institute (2013). *Marine fishing gears and fishing methods in South China Sea*. Beijing, China: China Ocean Press.
- Han, J., & Xu, H. (2004). *Overview of status and trend of trash fish from marine fisheries and their utilization, with special reference to aquaculture*. Qingdao, China: The Basic Status of Trash Fish in China.
- Hasan, M., & Halwart, M. (2009). Fish as feed inputs for aquaculture: Practices, sustainability and implications. *Practices, Sustainability and Implications*, 518, 407. 978-92-5-106419-1
- Hastings, A., Gaines, S. D., & Costello, C. (2017). Marine reserves solve an important bycatch problem in fisheries. *Proceedings of the National Academy of Sciences*, 114(34), 201705169. <https://doi.org/10.1073/pnas.1705169114>
- Huntington, T., & Hasan, M. (2009). Fish as feed inputs for aquaculture – Practices, sustainability and implications: A global synthesis. In M. R. Hasan, & M. Halwart (Eds.), *Fish as feed inputs for aquaculture: Practices, sustainability and implications* (pp. 1–61). Retrieved from <http://docs.niwa.co.nz/library/public/FAOfatp518.pdf>
- Jacobsen, N. S., Gislason, H., & Andersen, K. H. (2013). The consequences of balanced harvesting of fish communities. *Proceedings of the Royal Society B: Biological Sciences*, 281(1775), 1–9. <https://doi.org/10.1098/rspb.2013.2701>
- Jin, X. S., Shan, X. J., Li, X. S., Wang, J., Cui, Y., & Zuo, T. (2013). Long-term changes in the fishery ecosystem structure of Laizhou Bay. *China. Science China Earth Sciences*, 56(3), 366–374. <https://doi.org/10.1007/s11430-012-4528-7>
- Leadbitter, D. (2013). A risk based approach for promoting management regimes for trawl fisheries in South East Asia. *Asia Fisheries Science*, 26, 65–78.
- Liang, C., & Pauly, D. (2017). Growth and mortality of exploited fishes in China's coastal seas and their uses for yield-per-recruit analyses. *Journal of Applied Ichthyology*, 33(4), 746–756. <https://doi.org/10.1111/jai.13379>
- Liu, M., & Sadovy de Mitcheson, Y. (2008). Profile of a fishery collapse: Why mariculture failed to save the large yellow croaker. *Fish and Fisheries*, 9(3), 219–242. <https://doi.org/10.1111/j.1467-2979.2008.00278.x>
- Lundgren, R., Staples, D. J., Funge-Smith, S. J., & Clausen, J. (2006). *Status and potential of fisheries and aquaculture in Asia and the Pacific 2006*. Retrieved from <ftp://ftp.fao.org/docrep/fao/009/ag110e/ag110e00.pdf>
- Mallory, T. G. (2016). Fisheries subsidies in China: Quantitative and qualitative assessment of policy coherence and effectiveness. *Marine Policy*, 68, 74–82. <https://doi.org/10.1016/j.marpol.2016.01.028>
- Miao, W., & Liao, M. (2007). Analysis of feeds and fertilizers for sustainable aquaculture development in China. *FAO*, 497, 141–190.
- MOA (2017a). *China fishery statistical yearbook 2017 (1st ed.)*. Beijing, China: China Agriculture Press.
- MOA (2017b). *List of commercially important aquatic animals and plants under special state protection for resources*. Retrieved from [http://www.moa.gov.cn/sydw/stzz/zzyy/zzbh/201706/t20170608\\_5665914.htm](http://www.moa.gov.cn/sydw/stzz/zzyy/zzbh/201706/t20170608_5665914.htm)
- MOA (2017c). *Standard of catchable size of important fisheries species Part 1: Economic marine species*. Retrieved from [http://www.moa.gov.cn/govpublic/YYJ/201704/t20170401\\_5548770.htm](http://www.moa.gov.cn/govpublic/YYJ/201704/t20170401_5548770.htm)
- Naylor, R. L., Goldburg, R. J., Primavera, J. H., Kautsky, N., Beveridge, M. C. M., Clay, J., ... Troell, M. (2000). Effect of aquaculture on world fish supplies. *Nature*, 405(6790), 1017–1024. <https://doi.org/10.1038/35016500>
- Nguyen, T. T., Le, L. T., Tran, D. T., Cao, Q. L., Nguyen, H. T., Phan, D. N., & Tran, T. X. (2009). *Project on development planning for aquaculture sector in the Mekong Delta up to 2015 and strategic planning up to 2020 (in Vietnamese)*. Ho Chi Minh, Vietnam: Ministry of Agriculture & Rural Development.
- Parker, R. W. R., & Tyedmers, P. H. (2011). Uncertainty and natural variability in the ecological footprint of fisheries: A case study of reduction fisheries for meal and oil. *Ecological Indicators*, 16, 76–83. <https://doi.org/10.1016/j.ecolind.2011.06.015>

- Pauly, D., Belhabib, D., Blomeyer, R., Cheung, W. W. W. L., Cisneros-Montemayor, A. M., Copeland, D., ... Zeller, D. (2014). China's distant-water fisheries in the 21st century. *Fish and Fisheries*, 15(3), 474–488. <https://doi.org/10.1111/faf.12032>
- Pauly, D., Christensen, V., Dalsgaard, J., Froese, R., & Torres, F. J. (1998). Fishing down marine food webs. *Science*, 279, 860–863. <https://doi.org/10.1126/science.279.5352.860>
- Pikitch, E., Boersma, P. D. D., Boyd, I. L. I. L., Conover, D. D. O., Cury, P., Essington, T., ... Steneck, R. S. R. (2012). Little Fish, Big Impact: Managing a Crucial Link in Ocean Food Webs. Lenfest Ocean Program. Washington, DC. (108 pp.). Retrieved from [www.lenfestoceano.org/sites/default/files/online\\_ffft\\_summary.pdf](http://www.lenfestoceano.org/sites/default/files/online_ffft_summary.pdf)
- Pinsky, M. L., Jensen, O. P., Ricard, D., & Palumbi, S. R. (2011). Unexpected patterns of fisheries collapse in the world's oceans. *Proceedings of the National Academy of Sciences*, 108(20), 8317–8322. <https://doi.org/10.1073/pnas.1015313108>
- Qiu, Y. (2002). The situation of fishery resources in the northern South China Sea and the countermeasures for rational utilization (in Chinese). In *Survey China's exclusive economic zone and continental shelf* (pp. 360–367). Beijing, China: China Ocean Press.
- Rindorf, A., Cardinale, M., Shephard, S., De Oliveira, J. A. A., Hjørleifsson, E., Kempf, A., ... Vinther, M. (2017). Fishing for MSY: Using “pretty good yield” ranges without impairing recruitment. *ICES Journal of Marine Science*, 74(2), 525–534. <https://doi.org/10.1093/icesjms/fsw111>
- Rindorf, A., Dichmont, C. M., Levin, P. S., Mace, P., Pascoe, S., Prelezo, R., ... Clausen, L. W. (2017). Food for thought: Pretty good multispecies yield. *ICES Journal of Marine Science*, 74(2), 475–486. <https://doi.org/10.1093/icesjms/fsw071>
- Roberts, C. M., O'Leary, B. C., McCauley, D. J., Cury, P. M., Duarte, C. M., Lubchenco, J., ... Castilla, J. C. (2017). Marine reserves can mitigate and promote adaptation to climate change. *Proceedings of the National Academy of Sciences*, 114(24), 6167–6175. <https://doi.org/10.1073/pnas.1701262114>
- Sainsbury, K. J. (1984). Optimal mesh size for tropical multispecies trawl fisheries. *ICES Journal of Marine Science*, 41(2), 129–139. <https://doi.org/10.1093/icesjms/41.2.129>
- Shan, X., Sun, P., Jin, X., Li, X., & Dai, F. (2013). Long-term changes in fish assemblage structure in the Yellow River Estuary ecosystem, China. *Marine and Coastal Fisheries*, 5(1), 65–78. <https://doi.org/10.1080/19425120.2013.768571>
- Shen, G., & Heino, M. (2014). An overview of marine fisheries management in China. *Marine Policy*, 44, 265–272. <https://doi.org/10.1016/j.marpol.2013.09.012>
- Shepherd, C. J., & Jackson, A. J. (2013). Global fishmeal and fish-oil supply: Inputs, outputs and markets. *Journal of Fish Biology*, 83(4), 1046–1066. <https://doi.org/10.1111/jfb.12224>
- Siriraksophon, S., Pangsorn, S., & Laong-manee, P. (2009). The surimi industry in Southeast Asia: Trend and demand for raw materials. *Fish for the People*, 7(2), 2–8.
- Smith, A. D. M., Brown, C. J., Bulman, C. M., Fulton, E. A., Johnson, P., Kaplan, I. C., ... Tam, J. (2011). Impacts of fishing low-trophic level species on marine ecosystems. *Science*, 1147(6046), 1147–1150. <https://doi.org/10.1126/science.1209395>
- Smith, J. (1991). The atlantic and gulf menhaden purse seine fisheries: Origins, harvesting technologies, biostatistical monitoring, recent trends in fisheries statistics, and forecasting. *Marine Fisheries Review*, 53(4), 28–41.
- Sun, Z. (2014). *The general theory of fishing gears in the Yellow sea and the Bohai sea*. Beijing, China: China Ocean Press.
- Szuwalski, C. S., Burgess, M. G., Costello, C., & Gaines, S. D. (2017). High fishery catches through trophic cascades in China. *Proceedings of the National Academy of Sciences*, 114(4), 717–721. <https://doi.org/10.1073/pnas.1612722114>
- Tacon, A. G. J., Metian, M., Turchini, G. M., & De Silva, S. S. (2010). Responsible aquaculture and trophic level implications to global fish supply. *Reviews in Fisheries Science*, 18(1), 94–105. <https://doi.org/10.1080/10641260903325680>
- Tsikliras, A. C., & Stergiou, K. I. (2014). Size at maturity of Mediterranean marine fishes. *Reviews in Fish Biology and Fisheries*, 24(1), 219–268. <https://doi.org/10.1007/s11160-013-9330-x>
- Tveteras, S., Paredes, C. E., & Peña-Torres, J. (2011). Individual vessel quotas in Peru: Stopping the race for anchovies. *Marine Resource Economics*, 26(580), 225–232. <https://doi.org/10.5950/0738-1360-26.3.225>
- Wang, C. (2017). Review of fishmeal market in 2016 and prospects for 2017 (in Chinese). *Feed China*, 03, 18–21.
- Wang, L., & Yu, G. (2012). *Marine fishing gears and fishing method and management in East China Sea*. Hanzhou, China: Zhejiang Science and Technology Publishing House.
- Watson, R., Alder, J., & Pauly, D. (2006). Fisheries for forage fish, 1950 to the present. In J. Alder, & D. Pauly (Eds.), *On the multiple uses forage fish: From ecosystems to markets* (Vol. 14, pp. 1–19). Retrieved from [http://www.ecomarres.com/downloads/Chpt1\\_14-3.pdf](http://www.ecomarres.com/downloads/Chpt1_14-3.pdf)
- Yu, X. (2003). Implementation of minimum mesh size standards, Protection and rational use of fishery resources (in Chinese). *China Fisheries*, 07, 74–77.
- Zeller, D., Cashion, T., Palomares, M., & Pauly, D. (2018). Global marine fisheries discards: A synthesis of reconstructed data. *Fish and Fisheries*, 19(1), 30–39. <https://doi.org/10.1111/faf.12233>
- Zhou, D. (2007). Quality safety for aquaculture products of China and its management. *Global Trade Conference on Aquaculture: 29–31 May 2007, Qingdao, China*, 9, 213. Retrieved from <http://www.fao.org/3/a1454e/a1454e00.htm>
- Zhu, L., & Huang, S. (2014). The impact of fishery fuel subsidy policy on fishery resources and policy suggestions (in Chinese). *Journal of Shanghai Ocean University*, 23(04), 618–622.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Zhang W, Liu M, Sadovy de Mitcheson Y, et al. Fishing for feed in China: Facts, impacts and implications. *Fish Fish*. 2020;21:47–62. <https://doi.org/10.1111/faf.12414>